

Product datasheet for **MG216649**

Ffar2 (NM_001168512) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Ffar2 (NM_001168512) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Ffar2
Synonyms: GPCR43; Gpr43
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG216649 representing NM_001168512
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGACCCAGACTGGCACAGTTCCTTGATCCTCACGGCCTACATCCTCATCTTTCTTACTGGGCTCCCTG
CCAACCTGCTGGCCCTGCGGGCCTTCATGGGCCGGGTTCCGCCAGCCTCAGCCTGCCCTGTGCACATCCT
CCTGCTTAATCTGACCCTGGCGGACTTGCTCCTGTTGCTGCTGCTGCCCTCCGGATCGTGAAGCAGCA
TCCAACCTCCGCTGGTACCTACCAAAGATCGTGTGCGCGCTGACAGGCTTCGGCTTCTACAGCAGCATCT
ACTGCAGCACGTGGCTGCTGGCGGCATCAGCATGGAACGCTACCTGGGAGTGGCCTTCCGGTGCAGTA
CAAGTTATCCCGCCGGCCACTGTATGGAGTGATCGCTGCTCTGGTGGCCTGGATCATGTCCTTTGGCCAC
TGACCATCGTCATCATCGTTACAGTACCTGAACTCAACTGAGCAGGTGGGCACTGAGAACCAAATAACCT
GCTACGAGAACTTCAACCAAGAGCAGCTGGATGTGGTACTGCCCGTACGACTGGAGCTGTGCCTGGTCTT
GTTTTTCGTTCCCATGGCAGTCACCATCTTCTGTTATTGGCGCTTCGTGTGGATCATGCTCACGCAGCCC
CACGTTGGGGCTCAGAGGCGACGCCGGGAGTGGGCTGGCTGTTGTGACGCTTCTTAATTTCTGGTGT
GCTTTGGACCCTACAACATGTCCACCTGGTGGGTTCTACCTGAGGCAGAGCCCTCGTGGCGGGTGA
GGCTGTGGTGTTCAGTTCCTCAATGCCAGCCTGGATCCATTATTGTTCTACTTCTCCTCCTCCGTGGTG
CGCAGAGCTTTTGGAAAGGTTTGTACTGATCCGCAATCCTGCCTCCTATGCTGGGCAGGGGAGCCA
AAGAGACAGTGGAGGGGACCAAGATGGACAGGGGTGGAAGTCAAGCAGAAGGGGTACAGAGTTCTGAATT
TGTACCCGAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG216649 representing NM_001168512
 Red=Cloning site Green=Tags(s)

MTPDWHSSLILTAYILIFLTGLPANLLALRAFMRVRRQPAPVHILLNLTADLLLLLLLPFRIVEAA
 SNFRWYLPKIVCALTGFGFYSSICYSTWLLAGISMERYLGVAFPVQYKLSRRPLYGVIAALVAVIMSF
 GHCTIVIIQYLNSTEQVGTENQITCYENFTQEQLDVVLPVRLCLVLPVMAVTIFCYWRFVWIMLTQP
 HVGAQRRRRRAVGLAVVTLNLFVCFGPYNMSHLVGFYLRQSPSWRVEAVVSSLNASLDPLLFYFSSSVV
 RRAF GKLLLLIRNPASSMLGRGAKETVEGTMKDRGGSQAEGVQSSEFVTE

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_001168512

ORF Size: 990 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001168512.1 , NP_001161984.1
RefSeq Size:	2059 bp
RefSeq ORF:	993 bp
Locus ID:	233079
UniProt ID:	Q8VCK6
Cytogenetics:	7 B1

Gene Summary:

G protein-coupled receptor that is activated by a major product of dietary fiber digestion, the short chain fatty acids (SCFAs), and that plays a role in the regulation of whole-body energy homeostasis and in intestinal immunity. In omnivorous mammals, the short chain fatty acids acetate, propionate and butyrate are produced primarily by the gut microbiome that metabolizes dietary fibers. SCFAs serve as a source of energy but also act as signaling molecules. That G protein-coupled receptor is probably coupled to the pertussis toxin-sensitive, G(i/o)-alpha family of G proteins but also to the Gq family (PubMed:23589301). Its activation results in the formation of inositol 1,4,5-trisphosphate, the mobilization of intracellular calcium, the phosphorylation of the MAPK3/ERK1 and MAPK1/ERK2 kinases and the inhibition of intracellular cAMP accumulation. May play a role in glucose homeostasis by regulating the secretion of GLP-1, in response to short-chain fatty acids accumulating in the intestine (PubMed:22190648, PubMed:23589301). May also regulate the production of LEP/Leptin, a hormone acting on the central nervous system to inhibit food intake (PubMed:20399779). Finally, may also regulate whole-body energy homeostasis through adipogenesis regulating both differentiation and lipid storage of adipocytes (PubMed:16123168, PubMed:23589301). In parallel to its role in energy homeostasis, may also mediate the activation of the inflammatory and immune responses by SCFA in the intestine, regulating the rapid production of chemokines and cytokines (PubMed:23665276). May also play a role in the resolution of the inflammatory response and control chemotaxis in neutrophils (PubMed:19917676, PubMed:19865172). In addition to SCFAs, may also be activated by the extracellular lectin FCN1 in a process leading to activation of monocytes and inducing the secretion of interleukin-8/IL-8 in response to the presence of microbes. [UniProtKB/Swiss-Prot Function]