

Product datasheet for **MG216969**

Upf1 (NM_001122829) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Upf1 (NM_001122829) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Upf1
Synonyms:	B430202H16Rik; NORF1; PNORF-1; Rent1; Upflp
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG216969 representing NM_001122829 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAGTGTGGAGGCGTACGGCCCCAGCTCGAAACACTCACCTTCTTGGACACTGAGGAGGCCGAGCTGC
TCGGCGCCGACACCCAGGGCTCCGAGTTCGAATTCACCGACTTACCCTTCCCAGCCAGACGCAGACGCC
CCCCGGCGGCCCGGGCGCGGGAGGCCCGGGCGGAGCGGGCGCAGGCGCGCGGCCGCGCCAGCTCGAC
GCACAAGTTGGACCAGAGGGCATCTTGCAAATGGGGCTGTGGATGACAGTGTGGCAAGACCAGCCAGC
TGCTAGCTGAGCTGAACCTCGAGGAAGATGAAGAGGACACATACTACACTAAGGACTCCCAGTCCACGC
CTGCAGTACTGTGGAATCCATGATCCTGCCTGCGTGGTTTACTGTAATACCAGCAAGAAGTGGTTCTGC
AATGGCCGAGGAAATACTTCTGGCAGCCACATTGTGAATCACCTCGTGAGGGCAAATGCAAGGAAGTGA
CGCTGCACAAGGACGGGCCTCTGGGCGAGACCGTGTGGAGTGCTACAACCTGTGGCTGCCCAACGTCTT
CCTGCTGGGCTTCATCCCTGCGAAGGCCGACTCTGTGGTGGTGTGTTGTGCAGGCAGCCCTGTGCCAGC
CAGAGCAGCCTGAAGGACATCAACTGGGACAGCTCACAGTGGCAGCCCCTAATCCAGGACCGGTGCTTTC
TGTCATGGCTGGTCAAGATTCCTCTGAGCAGGAGCAGCTGCGAGCACGGCAGATCACGGCACAGCAGAT
CAACAAGCTGGAAGAGCTCTGGAAGGAAAATCCTTCAGCCACTCTGGAGGACCTGGAGAAGCCAGGCGTA
GACGAGGAGCCACAGCAGTGTCTCTGCGTTACGAGGATGCTTACCAGTACCAGAACATCTCGGGCCAC
TGGTCAAGCTGGAGGCTGACTATGACAAGAAGTTGAAGGAGTACAGACTCAAGATAACATCACGGTCAG
GTGGGACCTGGGCCTTAACAAGAAGAGAATCGCCTTCTTCACTTTGCCAAGACTGACTCTGGTAATGAG
GATTTAGTCATAATTTGGTTAAGAGACATGCGGCTCATGCAGGGTATGAGATCTGTCTGCGGTACAAAAG
GGGATCTGGCGCCCTGTGGAAGGGGATTGGCCACGTCATCAAGGTTCTGATAATTATGGTGATGAGAT
TGCTATTGAGCTCCGCAGCAGCGTGGGTGCCCTGTGGAAGTGACCCACAACCTCCAAGTGGATTTGTG
TGGAAGTCAACCTCTTTGATAGGATGCAGAGTCACTGAAGACCTTCGCTGTGGACGAGACCTCTGTGT
CAGGGTATATTTACCACAAGCTGCTGGGCCACGAGGTGGAGGATGTGGTATCAAGTCCAGCTGCCAAA
GCGCTTACAGCTCAGGGGCTCCCTGACCTCAACCACTCTCAGGTGTATGCTGTGAAGACCGTGTGCAG



[View online »](#)

AGACCACTCAGCCTCATCCAGGGCCCTCCAGGCACAGGCAAGACTGTGACATCAGCCACTATTGTCTACC
ACCTTGCTCGGCAGGGCAATGGGCCTGTACTGGTTTGTGCTCCAAGTAACATCGCTGTGGACCAGCTCAC
AGAGAAGATCCACCAGACAGGACTGAAGGTCGTACGCCTCTGTGCCAAGAGCCGTGAGGCCATTGACTCC
CCAGTGTCTTCCCTGGCTTTGCACAACCAGATCAGGAACATGGACAGCATGCCTGAGCTGCAGAAGCTGC
AGCAGCTAAAGGATGAGACAGGCGAGCTGTCATCTGCAGATGAGAAGCGGTACCGGGCGCTTAAGCGCAC
AGCTGAGAGAGAACTTCTCATGAATGCAGATGTCATATGCTGCACATGTGTGGTGTGCTGGTGACCCGAGG
CTGGCCAAAGATGCAGTTCCGTTCCATCCTTACGAGAGACACCCAGGCCACTGAGCTGAGTGCATGG
TGCTGTAGTCCCTGGGGCCAAGCAGCTAATCCTCGTTCGTTGACCACTGCCAGCTGGGCCAGTGGTGAT
GTGCAAGAAGGCAGCCAAGGCCGACTGTCACAATCGCTCTTTCGAGCGCTTGGTGGTGTGGGCATCCGG
CCCATCCGCCTGCAGGTGCAATACCGCATGCACCTGCACTCAGCGCCTTCCGTCACAACATCTTCTACG
AGGGCTCATTGCAGAATGGCGTCACTGCAGCGGATCGTGTCAAAAAGGCTTTGACTTCCAGTGGCCACA
ACCTGACAAGCCTATGTTCTTACGTGACGCAGGGCCAGGAGGATTGCCAGCTCTGGACATCTTAC
CTCAACAGGACGGAGGCCAATGTGGAGAAGATAACTACGAAGCTGTTGAAGGCAGGTGCAAAGCCTG
ACCAGATCGGCATCATCACCCCTACGAGGGCCAGCGCTTACTTGGTGCAGTACATGCAGTTCAGCGG
CTCCCTGCACAAAAGCTCTACCAGGAAGTGGAAATTGCCAGTGTGGACGCCTTCCAGGGCCGGGAGAAG
GACTTCATCATTCTGTCCCTGCGTGCAGGCAATGAACATCAGGGCATTGGGTTCCATAAGACCCCGGC
GTCTGAATGTGGCTCTCACAGAGCAAGATATGGCGTGATCATTGTGGTAACCCAAAGGCCCTGTGAA
GCAGCCCTGTGGAAATCACTGTGAGCTACTACAAGGAACAGAAGGCGTAGTGGAAAGGCCGCTCAAC
AACCTACGTGAGAGCCTCATGCAGTTACAGCAAGCCTCGCAAACCTGTCAACACTGTCAACCCGGGTGCC
GCTTACGACTACTGCCATGTACGATGCCCGTGAAGCCATCATCCCGGGTCTGTCTATGACCGCAGCAG
CCAGGGCCGGCCCTCGAATGTACTTCCAGACCCATGACCAGATTAGTATGATCAGCGCAGGCCCCAGC
CACGTGGCTGCCATGAACATCCCTATCCCTTCAACTTGGTTCATGCCTCCATGCCGCCACCTGGCTACT
TCGGACAGGCCAACGGGCCGAGCTGGTTCGGGCCACCCAAAACCAAGACTGGCCGTGGGGCCGGCCA
GAAGAACCCTTTGGGCTTCTGGGCCAGCCAGACCACCTTCCCAACAGCCAGGCCAGCCAGGAGCGTG
GCCTCCCAGCCCTTTTTCAGGGTGCCTCACACAGGGTTACGTGTCCATGAGCCAGCCCTCTCAGATGA
GCCAGCCTGGCCTCTCCCAGCCAGAAGTGTCCCAGGACAGCTACCTCGGTGATGAGTTTAAATCACAGAT
TGACGTGGCACTCTACAAGACTCCACATACCAGGGAGAGCGGGCATACCAGCACGGCCGGGTACCAGG
CTGTCCCAGTAC

ACGCGTACGCGGCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence:

>MG216969 representing NM_001122829

Red=Cloning site Green=Tags(s)

MSVEAYGPSSQTLTFLDTEEAELLGADTQGFSEFEFTDFTLPSQTQTPPGGPGGAGGPGGAGAGGAAGQLD
AQVGPEGILQNGAVDDSAKTSQLLAELNFEDEEDTYTKDLPVHACS YCGIHPACVVCNTSKKWF
NGRGNTSGSHIVNHLVRAKCKEVTLHKDGPLGETVLECYNGCRNVFLLGFIPAKADSVVLLCROPCAS
QSSLKIDINWSSQWQPLIQDRCFLSWLVKIPSEQEQLRARQITAQQINKLEELWKENPSATLEDLEKPGV
DEEPQHVLRLRYEDAYQYQNI FGPLVKLEADYDKLLESQTQDNITVRWDLGLNKKRIAFFTLPKTDSGNE
DLVIIWLRDMRLMQGDEICLRYKGDLAPLWKGIHVIVKVPDNYGDEIAIELRSSVGPVVEVTHNFQVDFV
WKSTSFDRMQSALKTFVAVDETSVSGYIYHKLLGHEVEDVVIKQQLPKRFTAQGLPDLNHSQVYAVKTVLQ
RPLSLIQPPGTGKTIVTSATIVYHLARQNGPVLVCAPSNI AVDQLTEKIHQTGLKVVRLCAKSREIDS
PVSFLALHNQIRNMDMPPELQKLQQLKDETGELSSADEKRYRALKRTAERELLMNADVICTCVGAGDPR
LAKMQFRSILIDESTQATEPECMVPVVLGAKQLILVGDHCQLGPVVMCKKAAGLSQSLFERL VVLGIR
PIRLQVQYRMHPALSAFPSNIFYEGSLQNGVTAADRVKKGDFQWPQPKPMFFYVTTQGGEEIASSGT
LNRTAANVEKITTLLKAGAKPDQIGIITPYEGQRSYLVQYMQFSGSLHTKLYQEVEIASVDAFQGREK
DFIILSCVRANEHQIGFLNDPRRLNVALTRARYGVII VGNPKALSKQPLWNHLLSYYKEQKALVEGPLN
NLRESLMQFSKPRKL VNTVNPGARFMTTAMYDAREAIIPGSVYDRSSQGRPSNMYYFQTHDQISMI
SAGPSHVAAMNIPIPFNLVMPMPPPPYPYGFQANGPAAGRGTGPKTKTGRGGRQKNRFGLPGPSQTTLPNSQASQDV
ASQPFSGAL TQGYVMSQPSQMSQPGLSQPELSQDSYLGDEFKSQIDVALSQDSTYQGERAYQHGGVTG
LSQY



TRTRPLE - GFP Tag - V


Restriction Sites:

SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:


ACCN: NM_001122829

ORF Size: 3372 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001122829.2](#), [NP_001116301.1](#)

RefSeq Size: 4631 bp

RefSeq ORF: 3375 bp

Locus ID: 19704

UniProt ID: [Q9EPU0](#)

Cytogenetics: 8 34.15 cM

Gene Summary:

RNA-dependent helicase and ATPase required for nonsense-mediated decay (NMD) of mRNAs containing premature stop codons. Is recruited to mRNAs upon translation termination and undergoes a cycle of phosphorylation and dephosphorylation; its phosphorylation appears to be a key step in NMD. Recruited by release factors to stalled ribosomes together with the SMG1C protein kinase complex to form the transient SURF (SMG1-UPF1-eRF1-eRF3) complex. In EJC-dependent NMD, the SURF complex associates with the exon junction complex (EJC) (located 50-55 or more nucleotides downstream from the termination codon) through UPF2 and allows the formation of an UPF1-UPF2-UPF3 surveillance complex which is believed to activate NMD. Phosphorylated UPF1 is recognized by EST1B/SMG5, SMG6 and SMG7 which are thought to provide a link to the mRNA degradation machinery involving exonucleolytic and endonucleolytic pathways, and to serve as adapters to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation. UPF1 can also activate NMD without UPF2 or UPF3, and in the absence of the NMD-enhancing downstream EJC indicative for alternative NMD pathways. Plays a role in replication-dependent histone mRNA degradation at the end of phase S; the function is independent of UPF2. For the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed. The ATPase activity of UPF1 is required for disassembly of mRNPs undergoing NMD (By similarity). Essential for embryonic viability.[UniProtKB/Swiss-Prot Function]