

Product datasheet for **MG219089**

Fbxo6 (NM_015797) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Fbxo6 (NM_015797) Mouse Tagged ORF Clone
Tag: TurboGFP
Symbol: Fbxo6
Synonyms: AA408845; FBG2; Fbs2; Fbx6b; Fbxo6b
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >MG219089 representing NM_015797
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGTCCACATCAACGAGCTGCCAGAGAACATTCTCCTGGAGCTGTTTCATCCATATCCCGCCCCACAGC
 TGCTGCGCAACTGCCGCTGGTCTGCCGCTCTGGCGAGACCTCATCGATGTGGTGTCCCTATGGAAGCG
 CAAGAGTCTTCGAGAGGGCTTCTCACCAAGACCGGTGCGAGCCGTGGAAGACTGGAAGTCTTCTAT
 ATCCTGTGCAGCCTGCAGAGAACCTCCTTCGGAACCCGTGTGCTGAAGAGAACCTGAGCTCATGGCGGA
 TAGACTCCAACGGAGGGGATCGCTGGAAGTGGAGACGCTCCCTGGGAGCTGTGGACAAGCTTTCTGA
 CAACAAGGTCAAGAAGTATTTTGTACCTCTTTTGTGAGATGTGCCTCAAATCCCAGATGGTGGACCTCAA
 GCTGAGGGCTACTGCGAGGAGCTGATGGACACCTTTCCGGCTGACATTGTGGTTAAGGACTGGGTTGCC
 CCAGAGCAGACTGTGGCTGCACCTATCAACTCCGGGTACAGCTGGCCTCTGCGGACTACATTGTCTTGGC
 CTCTTTTGTAGCCTCCACCTGTGACATCCAACAGTGGAATGATGCCAAATGGCAAGAGATTTCCACACC
 TTCTCTGATTACCCTCCAGGTGTCCGTACATCCTTTTCAACACGGGGGCCAGGACACTCAGTTCTGGA
 AAGGCTGGTACGGCCCCGTGTACCAACAGCAGCATCATTATCAGCCACAGGACAGCCAAGAACCCTCC
 CCCTGCCAGAACTCTACCGGAAGAACTGTAGTAATCGGAAGGAGACGGCGAGCTTCGGACTCCAACACT
 CATGAGGGTTTCTTCTGGCAAGGGCTATGGCAAAGGCTAAGGCGT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG219089 representing NM_015797
 Red=Cloning site Green=Tags(s)

MVHINELPENILLELFIHIPAPQLLRNCRVCRLWRDLIDVVSLLWKRKSLREGFFTKDRCEPVEDWKVFI
 ILCSLQRNLLRNPCAENLSSWRIDSNNGDRWKVETLPGSCGTSFPDNKVKKYFVTSFEMCLKSQMVDLK
 AEGYCEELMDTFRPDIVVKDWWAPRADCGCTYQLRVQLASADYIVLASFEPPTVFQQWVNDKQWEISHT
 FSDYPPGVRHILFQHGQDTQFWKGWYGPRVTNSSIIISHRTAKNPPPARTLPEETVVIGRRRRASDSNT
 HEGFFWQGLWQLRRR

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_015797

ORF Size: 885 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_015797.4 , NP_056612.1
RefSeq Size:	1955 bp
RefSeq ORF:	888 bp
Locus ID:	50762
UniProt ID:	Q9QZN4
Cytogenetics:	4 78.67 cM
Gene Summary:	Substrate-recognition component of some SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complexes. Involved in DNA damage response by specifically recognizing activated CHEK1 (phosphorylated on 'Ser-345'), promoting its ubiquitination and degradation. Ubiquitination of CHEK1 is required to insure that activated CHEK1 does not accumulate as cells progress through S phase, or when replication forks encounter transient impediments during normal DNA replication (By similarity). Involved in endoplasmic reticulum-associated degradation pathway (ERAD) for misfolded luminal proteins by recognizing and binding sugar chains on unfolded glycoproteins that are retrotranslocated into the cytosol and promoting their ubiquitination and subsequent degradation. Able to recognize and bind denatured glycoproteins, which are modified with not only high-mannose but also complex-type oligosaccharides. Also recognizes sulfated glycans.[UniProtKB/Swiss-Prot Function]