

Product datasheet for **MG219598**

Cd84 (NM_013489) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cd84 (NM_013489) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Cd84
Synonyms:	A130013D22Rik; CDw84; SLAMF5
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG219598 representing NM_013489 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCCAGCGCCATCTGTGGATCTGGTTCCTTTGCCTACAAACCTGGTCTGAAGCAGCAGGAAAAGATG
CAGACCCGATGGTAATGAATGGGATTCTGGGGAGTCAGTTACTTTCTCTTAAATATTCAAGAACAAA
GAAAATTGACAACATTGCCTGGACTTCTCAATCATCTGTTGCTTTATAAAACCAGGAGTCAATAAAGCT
GAAGTTACCATAACCCAGGGCACTTATAAAGGACGAATAGAAATCATAGATCAGAAGTATGACCTGGTCA
TTAGAGACCTGAGGATGGAAGATGCAGGAACCTTACAAAGCAGACATCAATGAAGAGAATGAGGAAACCAT
CACCAAGATCTACTACCTTCATATCTACCGTCGACTTAAAACACCAAAAATTACACAGAGTTTGATATCA
TCTTTGAACAATACCTGTAATATCACACTGACATGCTCTGTGGAAAAGGAAGAAAAGGATGTCACATATA
GCTGGAGTCCCTTTGGAGAGAAAAGCAATGTCCTTCAAATCGTCCACTCCCCATGGACCAAAAACCTGAC
CTACACATGTACAGCCCAGAACCCTGTCAGCAACAGTTCTGACTGTCTACTGTCCAGCAGCCATGTACA
GACTCCAAGCTTCCATCCTCGCCATGCTGTGTTGCCAGGAGGATTGGCCGTGCTTTCTGCTTATTC
TCATTCCGATGTTGGCATTCTGTTCCGTTTGTACAAGAGAAGGCGAGACAGGATTGCTCTGGAAGCAGA
TGATGTCTCAAAGAAAACAGTATATGCTGTAGTTTCAAGAAATGCTCAACCCACAGAGTCCAGAATCTAT
GATGAAATCCCTCAGTCCAAGATGCTGTCTGTAAAGAAAGATCCGGTGACCACCATTTATTCCTCAGTGC
AGCTTTCTGAGAAGATGAAGGAAACCAACATGAAGGACAGAAGTGCCTAAGGCTTTGGTAATGAAAT
TGTTGTC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG219598 representing NM_013489
 Red=Cloning site Green=Tags(s)

MAQRHLWIWFLCLQTWSEAAGKDADPMVMNGILGESVTFLLNIQEPKKIDNIAWTSQSSVAFIKPGVNKA
 EVTITQGTYKGRIEIIDQKYDLVIRDLRMEAGTYKADINEENEETITKIYYLHIYRRLKTPKITQSLIS
 SLNNTCNITLTCSVEKEEKDVITYSWSPFGEKSNVLQIVHSPMDQKLTYTCTAQNPVSNSSDSVTVQQPCT
 DTPSFHPRHAVLPGGLAVLFLILIPMLAFLFRLYKRRRDRIVLEADDVSKKTVYAVVSRNAQPTESRIY
 DEIPQSKMLSCKKDPVTIYSSVQLSEMKETNMKDRSLPKALGNEIVV

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_013489

ORF Size: 987 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_013489.2 , NP_038517.1
RefSeq Size:	3190 bp
RefSeq ORF:	990 bp
Locus ID:	12523
UniProt ID:	Q18PI6
Cytogenetics:	1 79.54 cM
Gene Summary:	<p>Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2 (PubMed:20962259). Can mediate natural killer (NK) cell cytotoxicity dependent on SH2D1A and SH2D1B (PubMed:20962259). Increases proliferative responses of activated T-cells and SH2D1A/SAP does not seem to be required for this process. Homophilic interactions enhance interferon gamma/IFNG secretion in lymphocytes and induce platelet stimulation via a SH2D1A/SAP-dependent pathway. May serve as a marker for hematopoietic progenitor cells (By similarity). Required for a prolonged T-cell:B-cell contact, optimal T follicular helper function, and germinal center formation (PubMed:20153220). In germinal centers involved in maintaining B cell tolerance and in preventing autoimmunity (PubMed:25801429). In mast cells negatively regulates high affinity immunoglobulin epsilon receptor signaling; independent of SH2D1A and SH2D1B but implicating FES and PTPN6/SHP-1 (By similarity). In macrophages enhances LPS-induced MAPK phosphorylation and NF-kappaB activation and modulates LPS-induced cytokine secretion; involving ITSM 2 (PubMed:20628063). Positively regulates macroautophagy in primary dendritic cells via stabilization of IRF8; inhibits TRIM21-mediated proteasomal degradation of IRF8 (By similarity).[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MG219598