

## **Product datasheet for MG222619**

## Cd79a (NM\_007655) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Cd79a (NM\_007655) Mouse Tagged ORF Clone

Tag: TurboGFP

Symbol: Cd79a

Synonyms: Ig-alpha; Iga; Igalpha; Ly-54; Ly54; mb-1

Mammalian Cell Neomycin

Selection:

Vector:

pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG222619 representing NM\_007655

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCCAGGGGGTCTAGAAGCCCTCAGAGCCCTGCTCTCCTCTCTTCTTGTCATACGCCTGTTTGGGTC
CCGGATGCCAGGCCCTGCGGGTAGAAGGGGGTCCACCATCCCTGACGGTGAACTTGGGCGAGGAGGCCCG
CCTCACCTGTGAAAACAATGGCAGGAACCCTAATATCACATGGTGGTTCAGCCTTCAGTCTAACATCACA
TGGCCCCCAGTGCCCACTGGGTCCTGGCCAGGGTACCACAGGCCAGCTGTTCTTCCCCGAAGTAAACAAGA
ACCACAGGGGCTTGTACTGGTGCCAAGTGATAGAAAACAACATATTAAAACGCTCCTGTGGTACTTACCT
CCGCGTGCGCAATCCAGTCCCTAGGCCCTTCCTGGACATGGGGGAAGGTACCAAGAACCGCATCATCACA
GCAGAAGGGATCATCTTGCTGTTCTGTGCAGTGGTGCCAGGGACGCTGCTGCTATTCAGGAAACGGTGGC
AAAATGAGAAGTTTGGGGTGGACATGCCAGATGACTATGAAGATGAAAATCTCTATGAGGGCCTGAACCT
TGATGACTGTTCTATGATGAGGACATCTCCAGGGGACTCCAGGGCACCTACCAGGATGTGGGCAACCTC

CACATTGGAGATGCCCAGCTGGAAAAGCCA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Protein Sequence:** >MG222619 representing NM\_007655

Red=Cloning site Green=Tags(s)


MPGGLEALRALPLLLFLSYACLGPGCQALRVEGGPPSLTVNLGEEARLTCENNGRNPNITWWFSLQSNIT WPPVPLGPGQGTTGQLFFPEVNKNHRGLYWCQVIENNILKRSCGTYLRVRNPVPRPFLDMGEGTKNRIIT AEGIILLFCAVVPGTLLLFRKRWQNEKFGVDMPDDYEDENLYEGLNLDDCSMYEDISRGLQGTYQDVGNL HIGDAQLEKP


TRTRPLE - GFP Tag - V

**Restriction Sites:** 


Sgfl-Mlul

**Cloning Scheme:** 





## Plasmid Map:



**ACCN:** NM\_007655

ORF Size: 660 bp

## Cd79a (NM\_007655) Mouse Tagged ORF Clone - MG222619

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

P11911

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 007655.4

RefSeq Size:1327 bpRefSeq ORF:663 bpLocus ID:12518

Cytogenetics: 7 13.49 cM

**UniProt ID:** 

**Gene Summary:** Required in cooperation with CD79B for initiation of the signal transduction cascade activated

by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to

internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into

proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of

immature B-cells.[UniProtKB/Swiss-Prot Function]