

Product datasheet for **MG222730**

Epm2a (NM_010146) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Epm2a (NM_010146) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Epm2a
Synonyms:	Tcrbk)TG-BFlv; Tg(TcraK; TG-B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG222730 representing NM_010146 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCTCTTCCGCTTCGGCGTGGTGGTGCCGCCCGCGGTGGCCGGCGCTCGGCAGGAGCTGCTGCTCGCCG
GCTCGCGGCCGAGTTAGGGCGCTGGGAGCCGACGGCGCCGTGCGTCTCAGGCCCGGGTACCGCGGC
GGGCGCCGCTGCGCTGGCGCTGCAGGAGCCCGCCTGTGGCTCGCCGAGGTGGAGCTGGAGGCGTACGAG
GAGGCCGGCGGGCGGAGCCGGGCCGCTTGACACGTTCTGGTACAAGTTCCTGCAGCGCGAGCCTGGAG
GCGAGCTGCACTGGGAAGGAAATGGACCTACCATGACCGTTGCTGCACATATAATGAGGACAACCTGGT
GGATGGTGTATTGTCTCCAGTAGGACTGGATTGAGGCCACTGGGCACACCAATGAAATGAAGCAC
ACAACAGACTTCTATTTAATATTGCTGGCCACCAAGCCATGCACTATTCAAGAATTCTACCAAATATCT
GGCTGGGTAGCTGCCCTCGCCAACCTGGAACATGTGACCATCAAATGAAGCATGAACTGGGAGTTACAGC
TGTGATGAATTTCCAGACTGAATGGGATATCATCCAGAATTTCTCAGGCTGCAACCGCTACCCTGAACCC
ATGACTCCAGACACCATGATGAAGCTGTATAAGGAAGAAGGCTTGTCTCATCTGGATGCCCACTCCAG
ACATGAGCACTGAGGGCCGAGTGCAGATGCTGCCACAGGCTGTGTCTCCTGCACGCGCTTCTGGAGAA
TGGACACACGGTGTATGTCCACTGCAACGCTGGCGTGGGTGCTCCACAGCTGCAGTGTGCGGCTGGCTC
CACTATGTGATTGGCTGGAATCTGCGCAAGGTGCAGTACTTCATCATGGCCAAGAGGCCTGCGGTCTACA
TTGACGAGGACGCTTTGGCTCAAGCACAACAAGACTTTTCTCAGAAGTTCGGGAAGGTTCACTCTCCAT
ATGCGCTTTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >MG222730 representing NM_010146
 Red=Cloning site Green=Tags(s)

MLFRFGVVVPPAVAGARQELLLAGSRPELGRWEPHGAVRLRPAGTAAGAAALALQEPGLWLAEEVELEAYE
 EAGGAEPGRVDTFWYKFLQREPGGELHWEONGPHHDCCTYNEDNLVDGVYCLPVGHWIEATGHTNEMKH
 TTDFYFNIAGHQAMHYSRILPNIWLGSCPRQLEHVTIKIKHELGVAVMNFQTEWDIIQNSSGCNRYPEP
 MTPDPTMMKLYKEEGLSYIWMPTPDMSTTEGRVQMLPQAVCLLHALLENGHTVYVHNCNAGVGRSTAAVCGWL
 HYVIGWNLKRVQYFIMAKRPAVYIDEDALAAQQDFSQKFGKVVHSSICAL

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_010146

ORF Size: 990 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_010146.2](#), [NP_034276.2](#)

RefSeq Size: 1079 bp

RefSeq ORF: 993 bp


Locus ID: 13853

UniProt ID: [Q9WUA5](#)

Cytogenetics: 10 A1

Gene Summary: Plays an important role in preventing glycogen hyperphosphorylation and the formation of insoluble aggregates, via its activity as glycogen phosphatase, and by promoting the ubiquitination of proteins involved in glycogen metabolism via its interaction with the E3 ubiquitin ligase NHLRC1/malin (PubMed:18040046, PubMed:18852261, PubMed:19036738, PubMed:23663739, PubMed:24430976, PubMed:24068615). Dephosphorylates phosphotyrosine and synthetic substrates, such as para-nitrophenylphosphate (pNPP), and has low activity with phosphoserine and phosphothreonine substrates (in vitro) (PubMed:16971387, PubMed:24430976). Has also been shown to dephosphorylate MAPT (PubMed:19542233). Shows strong phosphatase activity towards complex carbohydrates in vitro, avoiding glycogen hyperphosphorylation which is associated with reduced branching and formation of insoluble aggregates (PubMed:18040046, PubMed:18852261, PubMed:23663739). Forms a complex with NHLRC1/malin and HSP70, which suppresses the cellular toxicity of misfolded proteins by promoting their degradation through the ubiquitin-proteasome system (UPS) (PubMed:19036738, PubMed:24068615). Acts as a scaffold protein to facilitate PPP1R3C/PTG ubiquitination by NHLRC1/malin. Also promotes proteasome-independent protein degradation through the macroautophagy pathway (PubMed:20453062). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG222730