

Product datasheet for **MG225529**

II17rc (NM_134159) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	II17rc (NM_134159) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	II17rc
Synonyms:	1110025H02Rik; Gm19850; IL-17RC; IL17-RC; IL17-RL; II17rl
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




ACCN:	NM_134159
ORF Size:	2022 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_134159.4 , NP_598920.3
RefSeq Size:	2272 bp
RefSeq ORF:	2025 bp
Locus ID:	171095
UniProt ID:	Q8K4C2
Cytogenetics:	6 E3
Gene Summary:	Receptor for IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RA (PubMed:17911633, PubMed:20554964). Receptor for the heterodimer formed by IL17A and IL17B as part of a heterodimeric complex with IL17RA (By similarity). Has also been shown to be the cognate receptor for IL17F and to bind IL17A with high affinity without the need for IL17RA (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG225529