

Product datasheet for **MG225769**

Atf2 (NM_009715) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Atf2 (NM_009715) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Atf2
Synonyms:	Atf-2; CRE-BP; Creb2; D18875; D130078H02Rik; mXBP; Tg(Gzma-Klra1)7Wum
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MG225769 representing NM_009715
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGCACTGCCCTGGGTGTGCCAGATCAGACTCCAACGCCAACAAAGATTCCTAAAAAAGTGAAGAAG
 TGGTTTTGTTCAATGAGTTGGCAAGTCCATTTGAAAATGAATTCAGAAGGCTTCCGAAGATGACATTAA
 AAAAATGCCTCTAGATTTGTCCCCTCTTGAACACCCATCATAAGAAGCAAAAATTGAGGAGCCTTCTGTT
 GTAGAAAACAACCTACCAGGACAGCCCTTACCTACCCCGAGTCGACTACCAGTGATGAAAAGGAAGTAC
 CATTGGCACAACCTGCACAGCCACATCAGCTATCGTTCGTCCAGCATCATTACAGGTTCCCAATGTGCT
 GCTCACAAGTTCTGACTCAAGTGAATTATTCAACAAGCAGTACCTTACCAACCTCAAGTACTGTAATC
 ACCCAGGCACCATCTCTAACAGGCCAATTGTTCTGTACCAGGCCCATTTCTCTTCTTACATCTTC
 CTAATGGACAAAACATGCCCGTTGCTATTCCTGCATCAATTACAAGTTCTAATGTGCATGTTCCAGCTGC
 AGTCCCCTTGTTCGGCCAGTCACCATGGTGCCTAGTGTCCAGGAATCCAGGCCCTTCTCTCTCTCAA
 CCAGTCCAGTCAGAAGCAAAAATGAGATTAAGCTGCTTTGACCCAGCAACACCCCTCCAGTTACCAATG
 GTGATACTGTAAGGCCATGGCAGTGGATTGGTTAGGACTCAGTCAGAAGAGTCTCGCCACAGTCCTT
 GCAGCAGCCAGCCACCTCCACTACAGAACTCCGGCTTCTCCAGCTCACACAACCTCTCAGACCCAAAAT
 ACAAGTGGCCGTGAAGAAGAGCAGCTAATGAAGATCCTGATGAGAAAAGGAGGAAGTTTCTAGAACGAA
 ATAGAGCAGCAGCTTCAAGATGCCGACAAAAAGGAAAGTGTGGGTTCACTCCTTAGAGAAGAAAGCAGA
 AGACTTGAGTTCACTAAATGGCCAGCTGCAGAGCGAAGTACCCTGCTGAGAAAAGTGAAGTGGCCAGCTG
 AAACAGCTTCTTCTGGCTCATAAAGATTGCCCTGTAAGTCCATGCAGAAGAAGTCTGGCTATCATACTG
 CTGATAAAGATGACAGTTCAGAAGACCTTCTGTGCCAAGCAGTCCACATACAGAAGCGATCCAGCACAG
 CTCTGTACAGCACATCAATGGAGTCAGTTCAACATCAAAAAGCAGAAGCTGTAGCCACTTCAGTCTCACC
 CAGATGGCGGACCAGACGACGGAGCCTGCACTTTCACAGATTGTCATGGCTCCTCCTCCAGGCACAGC
 CCTCAGGAAGT

ACGCGTACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence:

>MG225769 representing NM_009715
 Red=Cloning site Green=Tags(s)

MHCPWVWPQTPTRFLKNCEEVGLFNELASPFENEFKKASEDDIKKMPDLSPLATPIIRSKIEEPSV
 VETTHQDSPLPHPESTTSDEKEVPLAQTAAQPTSAIVRPASLQVNVLLTSSDSSVIIQQAVPSPTSSTVI
 TQAPSSNRPIVPVPGPFLLLLHLPNGQTMPVAIPASITSSNVHVPAAVPLVRPVTMVPSVPGIPGPSSPQ
 PVQSEAKMRLKAALTQQHPPVTNGDTVKGHGSLVRTQSESRPQSLQQPATSTTETPASPAHTTPQTQN
 TSGRRRAANEDPDEKRRKFLERNRAAASRCRQKRKVVWVQSLKKAEDLSSLNGQLQSEVTLLRNEVAQL
 KQLLLAHKDCPVTAMQKSGYHTADKDDSSDLVSPSSPHTEAIQHSSVSTSNGVSTSKAEAVATSVLT
 QMADQSTEPALSQIVMAPPSSQAQPSGS



TRTRPLE – GFP Tag – V

Restriction Sites:

SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:


Plasmid Map:

ACCN: NM_009715

ORF Size: 1341 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)
OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_009715.3, NP_033845.1</u>
RefSeq Size:	4197 bp
RefSeq ORF:	1344 bp
Locus ID:	11909
Cytogenetics:	2 C3
Gene Summary:	<p>Transcriptional activator which regulates the transcription of various genes, including those involved in anti-apoptosis, cell growth, and DNA damage response. Dependent on its binding partner, binds to CRE (cAMP response element) consensus sequences (5'-TGACGTCA-3') or to AP-1 (activator protein 1) consensus sequences (5'-TGACTCA-3'). In the nucleus, contributes to global transcription and the DNA damage response, in addition to specific transcriptional activities that are related to cell development, proliferation and death. In the cytoplasm, interacts with and perturbs HK1- and VDAC1-containing complexes at the mitochondrial outer membrane, thereby impairing mitochondrial membrane potential, inducing mitochondrial leakage and promoting cell death. The phosphorylated form (mediated by ATM) plays a role in the DNA damage response and is involved in the ionizing radiation (IR)-induced S phase checkpoint control and in the recruitment of the MRN complex into the IR-induced foci (IRIF). Exhibits histone acetyltransferase (HAT) activity which specifically acetylates histones H2B and H4 in vitro. In concert with CUL3 and RBX1, promotes the degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. Can elicit oncogenic or tumor suppressor activities depending on the tissue or cell type (By similarity).[UniProtKB/Swiss-Prot Function]</p>