

Product datasheet for **MG225827**

Pik3cd (NM_008840) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pik3cd (NM_008840) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Pik3cd
Synonyms:	2410099E07Rik; 2610208K16Rik; AW545373; p110delta
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG225827 representing NM_008840 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCCCCCTGGGGTGGACTGCCCATGGAGTTCTGGACCAAGAGGAGAGCCAGAGCGTGGTTGTTGACT
TCTTGCTGCCACAGGGTCTACTTGAACCTCCCGTGTCCCGCAATGCCAACCTCAGCACCATCAAGCA
GGTGTGTGGCACCCTGCACAGTATGAGCCACTTCCACATGCTCAGTGACCCCGAGGCCATGTGTTT
ACCTGTGTGAACCAGACGGCGGAGCAGCAGGAGTTGGAGGATGAGCAGCGGAGGCTGTGCGACATCCAGC
CCTTCTGCCCGTGTGCGCCTGGTGGCCGAGAGGGGGACCGCTGAAGAAGCTCATTAACTCTCAGAT
CAGCCTCCTCATTGGCAAAGGTCTCCATGAGTTTGATTCCCTGCGGGACCCGGAAGTAAACGACTTCCGC
ACTAAGATGCGCCAGTTTTGTGAAGAGGCTGCTGCTCACCGCCAGCAGCTGGGCTGGGTGGAATGGCTGC
AGTACAGCTTCCCCCTGCAGCTGGAGCCCTCAGCAAGGGGTTGGCGGGCCGGCTTATTGCGTGTGAGCA
CCGAGCCCTGCTGGTCAACGTGAAGTTCGAGGGCAGTGAGGAGAGCTTACCTTCCAGGTATCCACCAAG
GACATGCCCTGGCACTGATGGCCTGTGCCCTCCGAAAAAGGCCACAGTGTCCGGCAGCCTCTGGTGG
AGCAGCCTGAGGAATATGCCCTGCAGGTGAACGGGAGGCAGCAATACCTTACGGCAACTACCCGCTCTG
CCACTTTCAGTACATCTGCAGCTGCCTACACAGCGGGCTGACCCCTCATCTGACCATGGTCCACTCCTCC
TCCATCCTTGCTATGCGGGATGAGCAGAGCAATCCTGCCCCCAAGTACAGAAACCAGTGCACAAACCTC
CCCCGATCCCTGCCAAGAAGCCCTCCTCTGTGTCCCTGTGGTCCCTGGAACAGCCATTCTCCATTGAGCT
GATCGAGGGCCGAAAAGTGAATGCTGACGAGCGGATGAAGCTGGTTGTTGAGCCGGGCTCTTCCATGGC
AATGAGATGCTGTGCAAGACTGTGTCAAGCTCGGAGGTGAATGTATGCTCAGAGCCCGTGTGGAAGCAGC
GACTGGAGTTCGATATCAGCGTCTGTGACCTCCCGCGCATGGCTCGACTCTGTTTTGCTCTATGCCGT
CGTGGAGAAGGCTAAGAAGGCACGCTCCACAAAGAAGAAGTCTAAGAAGGCGGACTGCCCATCGCTTGG
GCCAACCTCATGTATTGACTACAAAGATCAGCTCAAGACGGGGAGCGCTGCCTCTACATGTGGCCCT
CTGTCCCAGATGAGAAGGGAGAGCTGCTGAATCCTGCGGGTACAGTGCAGGGGAACCCCAACACGGAGAG
TGCCGCTGCCCTGCTCATCTACCTGCTGAGGTGGCCCCCACCCTGTGACTTCCCGCTCTGGAGAAG



[View online »](#)

ATCCTGGAGCTGGGGCGTCACGGGGAGCGTGGGCGCATCACGGAGGAGGAGCAGCTGCAGCTGCGGGAGA
 TCCTGGAACGGCGGGGATCCGGGAACTGTACGAACATGAGAAGGACCTGGTGTGGAAGATGCGCCACGA
 AGTCCAGGAGCATTTCCAGAGGCGCTGGCCCGCTGCTGCTGGTACCACCAAGTGGAAATAACATGAGGAT
 GTGGCCAGATGCTCTATTTGCTGTGCTCCTGGCCGAGCTGCCTGTGCTGAGCGCCCTGGAACCTCTGG
 ACTTTAGCTTTCCCGACTGCTACGTGGGCTCCTTCGCCATCAAGTCCCTTCGGAAGCTGACGGACGATGA
 GCTCTTCCAGTACCTTCTGCAGTGGTCAAGTCTCAAATATGAGTCTACCTGGACTGCGAGCTGACC
 AAATCTTGTGGCCGAGCCCTGGTAACCGCAAGATCGGACACTTCTGTTCTGGCACCTCCGCTCTG
 AGATGCAGTACCATCAGTGGCTCTGCGGTTTGGTCTCATCATGGAAGCCTACTGCAGAGGCAGCACCCA
 CCACATGAAGGTCTGATGAAGCAGGGGAAGCACTGAGCAAGCTTAAGGCACTGAATGACTTTGTGAAG
 GTGAGTCCCAGAAGACCACCAAGCCCCAAACCAAGGAGATGATGCATATGTGCATGCGCCAGGAGACCT
 ACATGGAGGCCCTGTCCACCTGCAGTCTCCACTCGACCCAGCACCTGCTGGAGGAAGTCTGCAGTGT
 GGAGCAGTGCACCTTCATGGACTCCAAATGAAGCCCTGTGGATCATGTACAGCAGCGAGGAGGCGGGC
 AGTGTGGCAACGTGGGCATCATCTTAAAGAACGGGGATGACCTCGCCAGGACATGCTGACTCTGCAGA
 TGATCCAGCTCATGGACGCTCTGTGGAAGCAGGAGGGCCTGGACCTGAGGATGACGCCCTACGGCTGCCT
 CCCCACGGGGACCGCACAGGTCTCATCGAGTGGTCTCCACTCGGACACCATCGCAACATCCAGCTG
 AACAAAAGCAACATGGCGGCCACAGCTGCCTTCAACAAGGACGCCCTGCTCAACTGGCTCAAGTCCAAGA
 ACCCTGGGGAGGCCCTGGATCGGGCCATTGAGGAATTCACCCTCTCTGTGCTGGCTACTGTGTGGCCAC
 ATATGTGCTGGGCATCGGTGACCGGCACAGCGACAACATCATGATCAGAGAGAGTGGGCAGCTCTTCCAC
 ATTGATTTTGGCCACTTTCTGGGAACTTCAAGACCAAGTTTGAATCAACCGAGAGCGCGTCCCCTTCA
 TTCTCACCTACGACTTTGTCCACGTGATCCAGCAGGGGAAGACTAACACAGTGAGAAGTTTGAAGGTT
 CCGAGGCTACTGTGAACGAGCCTACACCATCCTGCGGCCACGGGCTGCTTTTCTCCATCTCTTCGCC
 CTGATGCGGGCCGAGGTCTGCCTGAGCTTAGTTGCTCCAAGATATCCAGTATCTCAAGGACTCTCTGG
 CACTGGGAAGACGGAGGAAGAGCGCTAAAGCACTTCCGGGTGAAGTTCAACGAAGCTCTCCGAGAAAG
 CTGGAAAACCAAAGTCAACTGGCTGGCGCAATATGTGTCCAAGGATAACCGACAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence:

>MG225827 representing NM_008840
 Red=Cloning site Green=Tags(s)

MPPGVDCPMEFWTKEESQSVVDFLLPTGVYLNFPVSRNANLSTIKQVLWHRAQYEPLFHMLSDPEAYVF
 TCVNQTAEQELEDQRRLCDIQPFLPVLRLVAREGDRVKKLINSQISLLIGKGLHEFDSLDPVNDFR
 TKMRQFCEEAHRQQLGWVWELQYSFPLQLEPSARGWRAGLLRVSNRALLVNVKFESEESFTFQVSTK
 DMPLALMACALRKKATVFRQPLVEQPEEYALQVNGRHEYLGNYPPLCHFQYICSLHSGLTPHLTMVHSS
 SILAMRDEQSNPAPQVQKPRAKPPPAPAKKPSVSLWSLEQPFSEIELIEGRKVNADERMMLVQAGLFHG
 NEMLCKTVSSSEVNVCSPEVWKQRLEFDISVCDLPRMARLFCALYAVVEKAKKARSTKKKSKKADCP
 ANLMLFDYKDLKTGERCLYMWPSVPDEKCELLNPAGTVRGNPNTESSAALVIYLPVAPHPVYFPALEK
 ILELGRHGERGRITEEEQLQLREILERRGSGELYEHEKDLVWKMREVEHFPEALARLLVTKWKNHED
 VAQMLYLCSWPELPVLSALELLDFSPDCYVGSFAIKSLRKLTDDELQYLLQLVQVLKYESYLDCELT
 KFLLGRALANRKGHFLFWHLRSEMHPVSVALRFGLIMEAYCRGSTHMKVLMKQGEALSKLKALNDFVK
 VSSQKTKTPQTKEMMHCMRQETMEALSHLQSPDPSTLLEEVCVSEQCTFMDSKMKPLWIMYSSEEAG
 SAGNVGIIFKNGDDLQDMLTLQMIQLMDVLWKQEGDLRMTPTYGCLPTGDRGTGLIEVVLHSDTIANIQL
 NKSNMAATAAFNKDALLNWLKSKNPGEALDRAIEEFTLSCAGYCVATYVVLGIDRHSNIMIRESGQLFH
 IDFGHFLGNFKTKFGINRERVPIILTYDFVHVIQGGKTNSEKFERFRGYCERAYTILRRHGLLFLHLFA
 LMRAAGLPELSCSKDIQYLKDSLALGKTEEEALKHFRVKFNEALRESWTKVNWLAHNVSKDNRQ

TRTRPLE - GFP Tag - V


Restriction Sites:

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_008840

ORF Size: 3132 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_008840.3 , NP_032866.2
RefSeq Size:	5102 bp
RefSeq ORF:	3135 bp
Locus ID:	18707
Cytogenetics:	4 E2
Gene Summary:	<p>Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Mediates immune responses. Plays a role in B-cell development, proliferation, migration, and function. Required for B-cell receptor (BCR) signaling. Mediates B-cell proliferation response to anti-IgM, anti-CD40 and IL4 stimulation. Promotes cytokine production in response to TLR4 and TLR9. Required for antibody class switch mediated by TLR9. Involved in the antigen presentation function of B-cells. Involved in B-cell chemotaxis in response to CXCL13 and sphingosine 1-phosphate (S1P). Required for proliferation, signaling and cytokine production of naive, effector and memory T-cells. Required for T-cell receptor (TCR) signaling. Mediates TCR signaling events at the immune synapse. Activation by TCR leads to antigen-dependent memory T-cell migration and retention to antigenic tissues. Together with PIK3CG participates in T-cell development. Contributes to T-helper cell expansion and differentiation. Required for T-cell migration mediated by homing receptors SELL/CD62L, CCR7 and S1PR1 and antigen dependent recruitment of T-cells. Together with PIK3CG is involved in natural killer (NK) cell development and migration towards the sites of inflammation. Participates in NK cell receptor activation. Have a role in NK cell maturation and cytokine production. Together with PIK3CG is involved in neutrophil chemotaxis and extravasation. Together with PIK3CG participates in neutrophil respiratory burst. Have important roles in mast-cell development and mast cell mediated allergic response. Involved in stem cell factor (SCF)-mediated proliferation, adhesion and migration. Required for allergen-IgE-induced degranulation and cytokine release. The lipid kinase activity is required for its biological function.[UniProtKB/Swiss-Prot Function]</p>