

Product datasheet for **MG225878**

Akt1s1 (NM_026270) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Akt1s1 (NM_026270) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Akt1s1
Synonyms:	1110012J22Rik; AI227026; Lobe; Lobel; PRAS40
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>MG225878 representing NM_026270 Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGC**

ATGGCGTCTGGGCGGCCAGAGGAAGTGTGGGAAGCCGTCGTGGGGCGGCCGAGCGCTTTCAGGCCCGCA
 CTGGCACAGAGCTGGTATTACTGACTGCAGCGCCACCGCCGCGCCCGCCCTGGACCTGTGCCTATGC
 CGCCCATGGCCGCGGAGCCCTGGCAGAGGCGGCCGACGCTGCCTCCACGACATCGCACAGGCGCACAGG
 GCTGCCACTGCCACCGACCTCCTGGTCCCCACCAGCACCACAGCCGCCAGCCCTGCTCCTAGTCCAC
 CACCTCGGCCAGCCCTGGCCAGGAGGATGAGGAGGAAGATGAGGACGAGCCCACTGAAACAGAGACATC
 TGGGGAGCGGCTGGGCGGTAGCGATAATGGAGGTCTCTTCATGATGGATGAGGATGCCACCCTCCAGGAC
 CTGCCCCCTTCTGCGAGTCAGACCCGAGAGCACAGACGACGGCAGCCTGAGCGAGGAGACGCCCGCCG
 GTCCACAGCCTGTCCCAGCCCCCGGCCACAGCCCTGCCTACCCAGCAGTATGCCAAGTCTCTGCCCGT
 GTCGGTGCCAGTGTGGCCTTCAAGGAGAAGAGGACAGAAGCCCGATCGTCAGATGAGGAGAATGGCCCG
 CCCTCCTCGCCGACCTAGACCGAATAGCGGCCAGCATGCGCGCGCTGGTGCTGCGGGAGGCTGAGGACA
 CCCAGGTCTTCGGGGATCTTCGCGGCCGCGGCTCAATACCAGCGACTTCCAGAAGCTGAAGCGAAATA
 T

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA


[View online »](#)

Protein Sequence: >MG225878 representing NM_026270
 Red=Cloning site Green=Tags(s)

MASGRPEELWEAVVGAERFQARTGTEL VLLTAAPPPPPRPGPCAYAAHGRGALAEARRCLHDIAQHR
 AATATRPPGPPAPQPPSPAPSPPPR PALAREDEEDEDEPTETETSGERLGGSDNGGLFMMDEDATLQD
 LPPFCESDPESTDGSLSEETPAGPTACPPATALPTQQYAKSLPVSVPVWAFKEKRTEARSSDEENG
 PSSPDLDRIAASMRALVLREAEDTQVFGDLPRPRLNTSDFQKLKRY

TRTRPLE – GFP Tag – V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_026270

ORF Size: 771 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_026270.4, NP_080546.1</u>
RefSeq Size:	1578 bp
RefSeq ORF:	774 bp
Locus ID:	67605
UniProt ID:	<u>Q9D1F4</u>
Cytogenetics:	7 B3
Gene Summary:	<p>Subunit of mTORC1, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potently activates the protein kinase activity of mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Within mTORC1, AKT1S1 negatively regulates mTOR activity in a manner that is dependent on its phosphorylation state and binding to 14-3-3. Inhibits RHEB-GTP-dependent mTORC1 activation. Substrate for AKT1 phosphorylation, but can also be activated by AKT1-independent mechanisms. May also play a role in nerve growth factor-mediated neuroprotection.[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MG225878