


Product datasheet for **MG226429**

Otub2 (NM_001177841) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: Otub2 (NM_001177841) Mouse Tagged ORF Clone
 Tag: TurboGFP
 Symbol: Otub2
 Synonyms: 2010015L18Rik; 4930586I02Rik; AI413508; AW557219; OTB2; OTU2
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-AC-GFP (PS100010)
 E. coli Selection: Ampicillin (100 ug/mL)
 Restriction Sites: SgfI-MluI
 Cloning Scheme:

Cloning sites used for ORF Shuttling:




ACCN: NM_001177841
 ORF Size: 969 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001177841.1 , NP_001171312.1
RefSeq Size:	3239 bp
RefSeq ORF:	972 bp
Locus ID:	68149
UniProt ID:	Q9CQX0
Cytogenetics:	12 E
Gene Summary:	Hydrolase that can remove conjugated ubiquitin from proteins in vitro and may therefore play an important regulatory role at the level of protein turnover by preventing degradation. Mediates deubiquitination of 'Lys-11-', 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains, with a preference for 'Lys-63'-linked polyubiquitin chains (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MG226429