

Product datasheet for **MG226803**

Usp7 (NM_001003918) Mouse Tagged ORF Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | Usp7 (NM_001003918) Mouse Tagged ORF Clone |
| Tag: | TurboGFP |
| Symbol: | Usp7 |
| Synonyms: | 2210010O09Rik; AA409944; AA617399; AU019296; AW548146; C80752; Hausp |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-AC-GFP (PS100010) |
| E. coli Selection: | Ampicillin (100 ug/mL) |
| ORF Nucleotide Sequence: | >MG226803 representing NM_001003918 Red=Cloning site Blue=ORF Green=Tags(s) |

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAACCAACAGCAGCAGCAGCAGCAGCAGCAGAAAGCGGGCGAGCAGCAGCTGAGCGAGCCCGAGGACA
TGGAGATGGAAGCTGGAGATACAGATGACCCACCAAGAATTACTCAGAACCCTGTCATCAATGGGAATGT
AACCTGAGTGATGGGCACAGCAACGCAGAAGAGGACATGGAGGATGACACCAGTTGGCGCTCCGAGGCA
ACCTTTCAGTTCAGTGTGAGCGCTTCAGCAGACTGAGTGAGTCGGTCCCTTAGCCCTCCGTGTTTGTGC
GAAATCTGCCATGGAAGATTATGGTGATGCCACGCTTTTATCCAGACAGACCACCAAAAAAGCGTAGG
ATTCTTTCAGTGCAATGCTGAATCTGACTCCACGTCGTGGTCTGCCACGCACAAGCAGTGCTGAAG
ATAATAAATTACAGGGATGATGACAAATCATTCAGCCGGCGAATCAGCCACTTGTCTTCCACAAGGAAA
ACGACTGGGGATTCTCTAATTTTCATGGCCTGGAGTGAAGTGACTGATCCTGAGAAAGGATTCATCGATGA
TGACAAAGTCACCTTTGAAGTTTTGTACAGGCAGATGCTCCCATGGAGTTGCGTGGGACTCAAAGAAG
CATACTGGCTATGTGGCTTAAAGAATCAGGGAGCAACGTGTTACATGAATAGCTTGCTGCAGACCTTGT
TTTTACAAATCAGTTACGAAAGGCTGTGTACATGATGCCAACAGAGGGCGATGATTCATCTAAAAGCGT
CCCTTTAGCATTACAAAGAGTGTCTATGAATTACAGCACAGCGATAAGCCCGTAGGAACAAAAAGCTC
ACAAAGTCATTTGGTGGGAACTTTAGATAGCTTCATGCAACACGATGTTCAAGAGCTTTGTCGAGTGT
TGCTTGATAATGTGGAAAATAAGATGAAAGGCACCTGTGTAGAGGGCACCATTCCAAAGTTATCCGAGG
CAAAATGGTGTCTATATTCAGTGCAAAGAGCTAGACTACCGTTCTGATAGGAGAGAAGATTATTATGAT
ATCCAGCTAAGTATAAAAGGAAAGAAAAACATATTTGAATCATTTGTGGATTATGTGGCAGTAGAACAAC
TAGACGGTGACAATAAATACGATGCTGGGGAGCATGGCCTGCAGGAGGCAGAGAAAGGTGTGAAATTCCT
AACATTGCCCCAGTGTTACACCTGCAACTTATGAGGTTTCATGTATGACCCTCAGACGGACCAAAACATC
AAGATCAATGACAGGTTTGTGTTCCAGAACAGTTACCACTTGTGAATTTTTGCAAAAAACAGACCCTA
AGGACCCTGCAAATTATTTCTCCATGCAGTCTTGGTTCACAGTGGAGATAATCATGGTGGACATTACGT
GGTTTACCTAAACCCAAAGGGATGGCAAATGGTGTAAAGTTCGATGATGACGTGGTATCCAGGTGACT



[View online >](#)

AAAGAAGAAGCCATTGAGCACAATTATGGGGGTCATGATGATGATCTGTCTGTTTCGACACTGCACAAATG
 CCTATATGTTAGTGTACATCAGGAATCAAAGCTAAGTGAAGTGTACAAGCTGTACCCGACCATGATAT
 TCCTCAGCAGTTGGTGGAAACGATTGCAAGAAGAGAAAAGGATCGAGGCTCAGAAGCGGAAGGAGCGGCAG
 GAAGCCCATCTCTACATGCAAGTGCAGATAGTTGCAGAGGACCAGTTTTGTGGCCACCAAGGAAATGACA
 TGTACGACGAGGAGAAAGTGAGGTACACTGTGTTCAAGGTTCTGAAGAACTCCTCCCTGGCCGAGTTTGT
 TCAGAGCCTCTCCAGACCATGGGGTTCCACAAGATCAGATTCGATTATGGCCTATGCAAGCAAGAAGT
 AATGGAACAAAGCGCCTGCAATGTTAGATAATGAAGCTGACGGCAATAAAAACAATGATGTAAGTGAAGT
 ATAATGAAAACCTTGGACAATATTCTGGAAACAGTTGATCCTGAGCTGGCTGCTAGTGGAGCAACATT
 GCCCAAGTTTGATAAAGATCATGATGTAATGTTATTTTTGAAGATGTATGATCCTAAAACCCGGAGCTTG
 AATTACTGTGGACATATCTACACCCCATATCCTGTAATACTGACTTGTCTCCCGTTATGTGTGACA
 GAGCAGGATTTATCCAGGATACTAGCCTTATCCTCTATGAGGAAGTTAAGCCGAATTTAACAGAGAGAAT
 TCAGGACTATGATGTCTCTTGATAAAGCCCTTGATGAGCTGATGGATGGTACATCATAGTGTTCAG
 AAGGATGACCCAGAAAACGATAACAGTGAAGTGCCTGCTAAAGAGTATTTAGAGATCTCTACCACC
 GGGTCGACGTCTCTTGTGACAAAACGATCCCAATGACCCTGGATTTGTGGTACATTATCAAATAG
 GATGAAGTATTTTCAAGTTGCAAAGACAGTTGCACAGAGGCTCAACACAGACCCCATGCTGCTGCAGTTT
 TTCAAGTCTCAAGTTATAGGGACGGCCCGGTAATCCTCTTAGGCATAATTACGAAGGACTTTAAGAG
 ATCTTCTACAATCTTCAAGCCAAGACAACCTAAGAACTTTACTATCAGCAACTAAGATGAAAATCAC
 AGACTTTGAGAACAGGAGAAGTTTTAAGTGTATATGGTTAAACAGCCAATTTAGGGAAGAGGAAATAACA
 CTATACCCAGACAAGCATGGCTGTGTCCGGGACCTGTTAGAAGAATGTAAGAAAGCCGTGGAGCTCGGTG
 ACAAGCATCAGGGAGACTTAGGCTGCTAGAAATTTGTGAGCTACAAAATTTGGTGTGCATCAAGAAGA
 TGAAGTATTAGAATGCTTATCTCTGCAACGAGTCGGACTTTTGAATAGAGGAAATACCTTTGGACCAG
 GTAGACATAGATAAGGAAAACGAGATGCTAATCACAGTGGCACATTTCCACAAAGAGGTGTTTGGGACGT
 TTGGAATTCATTTTTGCTAAGGATACACCAGGTTGAGCATTTCGAGAGGTTGATGAAGCGGATACAGAG
 CTTGCTGGACATCCAGGAGAAGGAGTTTGAAGTTTAAATTTGAATTGTCATGATGGGCGGCACACAG
 TACATCAATGAAGATGAGTATGAAGTAAATTTGAAGGACTTTGAGCCACACCTGTAACATGTCTCATC
 CTCGGCCTTGGCTAGGGCTTGACCACTTCAACAAAGCCCAAGAGGAGTCGCTACACTTACCTAGAAAA
 GGCAATTAATAATCCACAAC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence:

>MG226803 representing NM_001003918
 Red=Cloning site Green=Tags(s)

MNHQQQQQQKAGEQQLSEPEDMEMEAGDTPPRITQNPVINGNVTLSDGHSNAEEDMEDDTSWRSEA
 TFQFTVERFSRLSESVLSPPCFVRNLPWKIMVMPRFYPDRPHQKSVGFFLQCNAESDSTSWSCHAQAVLK
 IINRDRDDKSFRRISHLFFHKENDWGF SNFMAWSEVTDPEKGF IDDDKVTFEV VQADAPHGVAVDSSK
 HTGYVGLKNQGATCYMNSLLQLFFTNQLRKAVYMPTEGDDSSKSVPLALQRVFEYELQHSKDPVGTKKL
 TKSFGWETLDSFMQHDVQELCRVLLDNVENKMKGT CVEGTIPKLFKGMVSYIQCKDVDYRSDRREDDYD
 IQLSIKGGKNIFESFVDYVAVEQLDGDNKYDAGEHGLQEAKEKGVKFLTLPPVLHLQLMRFMYDPQTDQNI
 KINDRFEFPEQLPLDEFLQKTDPKDPANYILHAVLVHSGDNHGGHYVYVLYLNPKGDKWCKFDDDDVSRCT
 KEEAIEHNYGGHDDLSVRHCTNAYMLVYIRESKLSVLQAVTDHDI PQQLVERLQEEKRIEAQKRKERQ
 EAHL YMQVQIVAEDQFCGHQGNMYDEEKVRYTVFKVLKNSSLAEFVQSLSQTMGFPQDQIRLWPMQARS
 NGTKRPAML DNEADGNKTMIELSDNENPWTIFLETVDPELAASGATLPKFDKDHVMLFLKMYDPKTRSL
 NYCGHIYTPISCKIRDLLPVMCDRAGFIQDTSILYEEVKPNLTERIQDYDVS LDKALDELMDGDIIVFQ
 KDDPENDNSELPTAKEYFRDLYHRVDVIFCDKTI PNDPGFVVTL SNRMNYFQVAKTVAQRLNDPMLLQF
 FKSQGYRDGPGNPLRHNYEGLRDLLQFFKPRQPKKLYYQQLKMKITDFENRRSFKCIWLSQFRIIEEIPLDQ
 VYIDKENEMLITVAHFHKEVFGTFGIPFLLR IHQGEHFREVMKRIQSLLDIQEKEFEKFKFAIVMMGRHQ
 YINEDEYEVNLDKDFEPQPGNMSHPRPWLGLDHFNKAPKRSRYTYLEKAIKIHN


TRTRPLE - GFP Tag - V

Restriction Sites:


Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



Plasmid Map:



ACCN: NM_001003918
 ORF Size: 3309 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001003918.2](#), [NP_001003918.2](#)

RefSeq Size: 5427 bp

RefSeq ORF: 3312 bp

Locus ID: 252870

UniProt ID: [Q6A4J8](#)

Cytogenetics: 16 A1

Gene Summary:

Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN, KMT2E and DAXX (PubMed:21268065, PubMed:14719112, PubMed:19946331). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation. Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis. Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis. Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity. In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML. Deubiquitinates KMT2E preventing KMT2E proteasomal-mediated degradation (By similarity). Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV-induced degradation of ERCC6 (By similarity). Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1. Deubiquitinates alkylation repair enzyme ALKBH3. OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions (By similarity). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (By similarity). Able to mediate deubiquitination of histone H2B; it is however unsure whether this activity takes place in vivo (PubMed:27863226). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains. Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:23973222). Plays a role in the maintenance of the circadian clock periodicity via deubiquitination and stabilization of the CRY1 and CRY2 proteins (PubMed:27123980). Deubiquitinates REST, thereby stabilizing REST and promoting the maintenance of neural progenitor cells (By similarity).[UniProtKB/Swiss-Prot Function]