

Product datasheet for MG227171

Mapk3 (NM_011952) Mouse Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Expression Plasmids
Mapk3 (NM_011952) Mouse Tagged ORF Clone
TurboGFP
Mapk3
Erk-1; Erk1; Ert2; Esrk1; Mnk1; Mtap2k; p44; p44erk1; p44mapk; Prkm3
Neomycin
pCMV6-AC-GFP (PS100010)
Ampicillin (100 ug/mL)
<pre>>MG227171 representing NM_011952 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
ATGGCGGCGGCGGCGGCGGCGCGGGGGGGGGGGGGGGGG

TACCGAGCCCCAGAGATCATGCTTAATTCCAAGGGCTACACCAAATCCATCGACATCTGGTCTGTGGGCT GCATTCTGGCTGAGATGCTCTCCAACCGGCCCATCTTCCCCGGCAAGCACTACCTGGACCAGCTCAACCA CATTCTAGGTATCTTGGGTTCCCCATCCCAGGAGGACCTTAATTGCATCATTAACATGAAGGCCCGAAAC TACCTGCAGTCTCTGCCCTCGAAAACCAAGGTGGCTTGGGCCAAGCTCTTTCCTAAATCTGACTCCAAAG CTCTTGACCTGCTGGACCGGATGTTAACCTTCAACCCAAACAAGCGCATCACAGTAGAGGAAGCGCTGGC TCACCCTTACCTGGAACAGTACTACGATCCGACAGATGAGCCAGTGGCCGAGGAGCCATTCACCTTCGAC ATGGAGCTGGATGACCTCCCAAGGAGCGGCTGAAGGAGTTGATCTTCCAAGAGAACAGCCCGCTTCCAGC



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

CAGGGGCGCCAGAGGGCCCC



ACCN:	NM_011952
ORF Size:	1140 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE Mapk3 (NM_011952) Mouse Tagged ORF Clone – MG227171

Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 011952.2, NP 036082.1</u>
RefSeq Size:	1772 bp
RefSeq ORF:	1143 bp
Locus ID:	26417
UniProt ID:	<u>Q63844</u>
Cytogenetics:	7 69.25 cM
Gene Summary:	Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated

by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements. The MAPK/ERK cascade plays also a role in initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors. About 160 substrates have already been discovered for ERKs. Many of these substrates are localized in the nucleus, and seem to participate in the regulation of transcription upon stimulation. However, other substrates are found in the cytosol as well as in other cellular organelles, and those are responsible for processes such as translation, mitosis and apoptosis. Moreover, the MAPK/ERK cascade is also involved in the regulation of the endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC); as well as in the fragmentation of the Golgi apparatus during mitosis. The substrates include transcription factors (such as ATF2, BCL6, ELK1, ERF, FOS, HSF4 or SPZ1), cytoskeletal elements (such as CANX, CTTN, GJA1, MAP2, MAPT, PXN, SORBS3 or STMN1), regulators of apoptosis (such as BAD, BTG2, CASP9, DAPK1, IER3, MCL1 or PPARG), regulators of translation (such as EIF4EBP1) and a variety of other signaling-related molecules (like ARHGEF2, FRS2 or GRB10). Protein kinases (such as RAF1, RPS6KA1/RSK1, RPS6KA3/RSK2, RPS6KA2/RSK3, RPS6KA6/RSK4, SYK, MKNK1/MNK1, MKNK2/MNK2, RPS6KA5/MSK1, RPS6KA4/MSK2, MAPKAPK3 or MAPKAPK5) and phosphatases (such as DUSP1, DUSP4, DUSP6 or DUSP16) are other substrates which enable the propagation the MAPK/ERK signal to additional cytosolic and nuclear targets, thereby extending the specificity of the cascade.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Circular map for MG227171

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US