

Product datasheet for **MG227189**

Irak1 (NM_008363) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Irak1 (NM_008363) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Irak1
Synonyms:	AA408924; I11rak; IRAK; IRAK-1; IRAK1-S; IRAK1b; mPLK; Plpk
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MG227189 representing NM_008363
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCCGGGGGGCCGGGCCCGGGGAGCCTGTGGTTCCAGGCGCCAGCACTTCTTGTACGAGGTGCCAC
 CCTGGGTTATGTGCCGCTTCTACAAAGTATGGACGCCCTGGAGCCCGCGACTGGTGCCAGTTCGCGGC
 CTTGATCGTGCACGACCAGACAGAGCTGCGGCTGTGCGAGCGCTCCGAGCAGCGCACAGCCAGTGTCTG
 TGGCCCTGGATCAACCGCAACGCGCGCGTAGTGACCTCGTTCACATCCTCACGCACCTGCAGTGTCTG
 GTGCGAGGGACATCATCACAGCCTGGCACCTCCTGCCCCGTTGTGCCCAAGCACCCTGCCCAAG
 GCCCAGCAGCATCTGCAAGCTCTGAGGCCGGGACTGGAGCCCCGAAATTGCAGTCTCTGCCTCC
 ACCTTCTCTCCAGCTTTTCCAGGCTCCAGACCCATTCTGAGTCAGAGCTCCTCCAGTTCCACTCC
 CTGTTTCCCTCGGGCCACCACTACCATCTCAGCCCTTCTCCACCAAGTCAAGCCAGAGAGCCAGT
 GTCTGGCTCCAAGAGCCCATCCTCCCGTTTTGCTGGCCCTTCTGTGAGATTTCCAAGGCACCTGC
 AACTTCTCTGAAGAACTCAGATTGGAGAGGGTGGTTTTGGATGTGTGTACCGAGCAGTCATGAGAAATA
 CTACATATGCTGTGAAGAGACTGAAGGAGGAAGCTGACCTAGAGTGGACTATGGTGAAACAGAGTCTT
 AACAGAGGTGGAACAGCTATCAAGTTTTCGTACCCAAATATCGTAGACTTTGCTGGCTACTGTGCAGAG
 AGTGGCTTATACTGCCTTGTATGGCTTCTTGCCCAATGGCTCCTTGGAGGATCAGTCCACCTCAGA
 CCCAGGCTGCTCCCACTTTCTGGCCTCAACGACTGGACATTTCTTGGGCACAGCCCGGCTATTCA
 GTTTTTACATCAGGATAGCCCCAGCCTTATCCATGGAGACATCAAGATTCTAACGTGCTTCTGGATGAG
 AGACTGATGCCAAGCTGGGAGACTTTGGCCTGGCTCGTTTTCAGCCGCTTTCGGGGGCCAAAGCAAG
 AGAGCAGTACTGTGCCCGGACTTCCACAGTTCGAGGTACCCTCGCCTACTTGCCTGAGGATACATCAA
 GACAGGCCGACTGGCTGTGGACACCGATACCTTCAGCTTTGGGGTGGTAATACTGGAGACCCTTGCTGGT
 CAGAGGGCTGTGAGGACACAAGGTGCAAAGACCAAGTATTTGAAAGACCTGATTGAAGATGAGGCTGAAG
 AGGCTGGAGTGACCCTGAAAAGCACCCAGCCTACTCTGTGGTGGGTGTAGCCACGGATGCTTGGCTGC
 TCCAATTGCTGCCAGATCTATAAGAAGCACCTGGACTCCAGACCTGGGCCCTGCCACCCAGTTGGGC
 CTGGCCCTGGCTCAACTAGCTTGTGCTGCATGCACCGTCCGGCCAAGAAGAGGCCCCCATGACCCAGG
 TATAAAGAGACTAGAAGGGCTCAGGCAGGGCTCCCTGGGAGCTAGAGGTTGCCGGCCATGGCTCCCC
 TTCCCCACAGGAACTCCTACATGTCTACCCTGGCAGTCCCAGAGTGGGGATGAACCATGGCAGCCT
 CTAGTAGTGACCACAAGAGCCCCAGCCAGGCTGCCAGCACTCCAGAGAAGTCCAACAGCCAGTGG
 AAAGTGATGAGAGTGTCCCGGCCCTCTCTGCTACCCTGCATTCTGGCACTTGACTCCAGTTCCACCC
 AAGCCCAGCGTCTTCAGAGAGGCTAGCTGTACCCAGGGAGGCACTACCAGAGAATCAAGTGTGAGGAGT
 AGCCCAGGCTTCCAGCCTACAACCATGGAAGGCTCACCCACGGGCAGCTCATCCCTGCTGTGCATCAGAAC
 CACCACAGATCATCATCAACCCAGCCGACAGAAGATGGTACAAAAGCTGGCTCTTTATGAAGAAGGGGT
 CTTGGATAGCCTGCAACTGCTGTGCATCAGGCTTTTCCAGGCTTGGATTTAGAACCTGAAAAGAGCCAG
 GGACCTGAAGAAAGTGATGAATTCCAGAGC

ACGCGTACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence: >MG227189 representing NM_008363
 Red=Cloning site Green=Tags(s)

```
MAGGPGPGPEVVPVGAQHFLYEVPWVMCRFYKVMdalePADWCQFAALIVRDQTELRLCERSEQRTASVL
WPWINRNARVADLVHILTHLQLLRARDIITAWHPPAPVPPSTAAPRPSSIAGSEAGDWSPRKLQSSAS
TFLSPAFPGSQTHSESELLQVPLPVSLGPPLPSSAPSSTKSSPESPVSGLQRAHPSFPCWPFCEISQGTC
NFSEELRIGEGGFCVYRAVMRNTTYAVKRLKEEADLEWTMVKQSFLTEVEQLSRFRHPNIVDFAGYCAE
SGLYCLVYGFPLPNGSLEDQLHLQTQACSPLSWPQRDLILLGTARAIQFLHQDSPSLIHGDIKSSNVLLDE
RLMPKLGDFLARFSRFAGAKASQSSTVARTSTVRGTLAYLPEEYIKTGR LAVD TDF SFGV VILETLAG
QRAVRTQGAKT KYL KDL IEDEAEEAGVTLKSTQPTLWVG VATDAAWAPIAAQIYKHLDSRPGPCPPQLG
LALAQLACCCMHRRAKKRPPMTQVYKRLEGLQAGPPWELEVAGHGSPSPQENSYMSTTGSAGSGDEPWQP
LVVTTTRAPAQAQQLRSPNQPVESDESVPGLSATLHSHWLTGSHSPSPASFREASCTQGGTTRESSVRS
SPGFQPTTMEGSPGSSLLSSEPPQIIINPARQKMQVQLALYEEGVLDLSQLLSSGFFPGLDLEPEKSQ
GPEESDEFQS
```


TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



Plasmid Map:


ACCN: NM_008363

ORF Size: 2130 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_008363.2](#), [NP_032389.2](#)

RefSeq Size: 3831 bp

RefSeq ORF: 2133 bp

Locus ID: 16179

UniProt ID: [Q62406](#)

Cytogenetics: X 37.61 cM

Gene Summary: Serine/threonine-protein kinase that plays a critical role in initiating innate immune response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways. Is rapidly recruited by MYD88 to the receptor-signaling complex upon TLR activation. Association with MYD88 leads to IRAK1 phosphorylation by IRAK4 and subsequent autophosphorylation and kinase activation. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin-binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6 complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent degradation. Phosphorylates the interferon regulatory factor 7 (IRF7) to induce its activation and translocation to the nucleus, resulting in transcriptional activation of type I IFN genes, which drive the cell in an antiviral state. When sumoylated, translocates to the nucleus and phosphorylates STAT3 (By similarity). [UniProtKB/Swiss-Prot Function]