

## Product datasheet for **MG227319**

### Lck (NM\_001162432) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Lck (NM_001162432) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Lck
Synonyms:	Hck-3; Lsk; Lskt; p56; p56Lck
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>MG227319 representing NM\_001162432  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**GCGATCGCC**

ATGGGGCCTCTGAGCTGACGATCTCGGGATCATGGGCTGTGTCTGCAGCTCAAACCTGAAGATGACT  
 GGATGGAGAACATTGACGTGTGTGAAAACCTGCCACTATCCCATAGTCCCCTGGACAGCAAGATCTCGCT  
 GCCCATCCGGAATGGCTCTGAAGTGCGGGACCCACTGGTCACCTATGAGGGATCTCTCCACCAGCATCC  
 CCGCTGCAAGACAACCTGGTTATCGCCCTGCACAGTTATGAGCCCTCCCATGATGGAGACTTGGGCTTTG  
 AGAAGGGTGAACAGCTCCGAATCCTGGAGCAGAGCGGTGAGTGGTGAAGGCTCAGTCCCTGACGACTGG  
 CCAAGAAGGCTTCATTCCCTTCACTTCGTGGCGAAAAGCAAACAGCCTGGAGCCTGAACCTTGGTTCTTC  
 AAGAATCTGAGCCGTAAGGACGCCGAGCGGCAGCTTTTGGCGCCCGGAACACGCATGGATCCTTCTGA  
 TCCGGGAAAGCGAAAGCACTGCGGGTCTTTTCCCTGTCGGTACAGAGACTTCGACCAGAACCAGGGAGA  
 AGTGGTGAACATTACAAGATCCGTAACCTAGACAACGGTGGCTTCTACATCTCCCTCGTATCACTTTT  
 CCCGGATTGCACGATCTAGTCCGCCATTACACCAACGCCTCTGATGGGCTGTGCACAAAGTTGAGCCGTC  
 CTTGCCAGACCCAGAAGCCCCAGAAACCATGGTGGGAGGACGAATGGGAAGTTCCAGGGAAACACTGAA  
 GTTGGTGGAGCGGCTGGGAGCTGGCCAGTTCGGGGAAGTGTGGATGGGACTACAACGGACACAGGAAG  
 GTGGCGGTGAAGAGTCTGAAACAAGGGAGCATGTCCCGGACGCCTTCTGGCTGAGGCTAACCTCATGA  
 AGCAGCTGCAGCACCCGCGGCTAGTCCGGCTTTATGCAGTGGTCAACCCAGGAACCCATCTACATCATCAC  
 GGAATACATGGAGAACGGGAGCCTAGTAGATTTTCTCAAGACTCCCTCGGCATCAAGTTGAATGTCAAC  
 AAACCTTTTGGACATGGCAGCCCAGATTGCAGAGGGCATGGCGTTCATCGAAGAACAGAATTACATCCATC  
 GGGACCTGCGCGCCCAACATCCTGGTGTCTGACACGCTGAGCTGCAAGATTGCAGCTTTGGCTGGC  
 GCGCCTCATTGAGGACAATGAGTACACGGCCCGGAGGGGCCAAATTTCCCATTAAGTGGACAGCACCA  
 GAAGCCATTAACTATGGGACCTTACCATCAAGTACAGACGTGTGGTCTTCCGGGATCTTGCTTACAGAGA  
 TCGTCACCACGGTCAATCCCTTACCAGGAATGACCAACCCTGAAGTCATTGAGAACCCTGGAGAGAGG  
 CTACCGCATGGTGAACCTGACAACCTGTCGGAAGAGCTGTACCACCTCATGATGCTGTGCTGGAAGGAG  
 CGCCAGAGGACCGCCACGTTTGACTACCTCGGAGTGTCTGGATGACTTCTCACAGCCACAGAGG  
 GCCAGTACCAGCCCCAGCCT

**ACGCGTACGCGGCCGCTCGAG** - GFP Tag - GTTTAA

**Protein Sequence:**

>MG227319 representing NM\_001162432  
 Red=Cloning site Green=Tags(s)

MGASELTISGIMGVCSSNPEDDWMENIDVCENHYIIVPLDSKISLPIRNGSEVRDPLVTYEGSLPPAS  
 PLQDNLVIALHSYEPSHDGDLGFEKGEQLRILEQSGEWWKAQSLTTQEGFIPFNFVAKANSLEPEPWFF  
 KNLNRKDAERQLLAPGNTHGSFLIRESESTAGSFLSVRDFDQNGEVVKHYKIRNLDNGGFYISPRITF  
 PGLHDLVRHYTNASDGLCTKLSRPCQTQKPKPWWEDEWEVPRETLKLVRLGAGQFGEVWVMGYNGHTK  
 VAVKSLKQGSMPDAFLAEANLQKQHPRLVRLYAVVTQEPIYIITEYMENGLVDFLKTSPGKLVN  
 KLLDMAAQIAEGMAFIEEQNYIHRDLRAANILVSDTLCKIADFLARLIEDNEYTAREGAKFPIKWTAP  
 EAINYGTFTIKSDVWSFGILLTEIVTHGRIPYPGMTNPEVIQNLERGYRMRPDNCPPEELYHLMMLCWKE  
 RPEDRPTFDYLRSLDDFFTATEGYQYQP

**TRTRPLE** - GFP Tag - V

**Restriction Sites:**


Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



Plasmid Map:



<b>ACCN:</b>	NM_001162432
<b>ORF Size:</b>	1560 bp
<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001162432.1</a> , <a href="#">NP_001155904.1</a>
<b>RefSeq Size:</b>	2074 bp
<b>RefSeq ORF:</b>	1563 bp
<b>Locus ID:</b>	16818
<b>Cytogenetics:</b>	4 63.26 cM

**Gene Summary:**

Non-receptor tyrosine-protein kinase that plays an essential role in the selection and maturation of developing T-cells in the thymus and in the function of mature T-cells. Plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways. Constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors. Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, thereby recruiting the associated LCK protein to the vicinity of the TCR/CD3 complex. LCK then phosphorylates tyrosine residues within the immunoreceptor tyrosine-based activation motifs (ITAM) of the cytoplasmic tails of the TCR-gamma chains and CD3 subunits, initiating the TCR/CD3 signaling pathway. Once stimulated, the TCR recruits the tyrosine kinase ZAP70, that becomes phosphorylated and activated by LCK. Following this, a large number of signaling molecules are recruited, ultimately leading to lymphokine production. LCK also contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, which leads to hyperphosphorylation and activation of LCK. Also plays a role in the IL2 receptor-linked signaling pathway that controls the T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Phosphorylates other substrates including RUNX3, PTK2B/PYK2, the microtubule-associated protein MAPT, RHOH or TYROBP (By similarity). Interacts with UNC119; this interaction plays a crucial role in activation of LCK (By similarity).  
[UniProtKB/Swiss-Prot Function]