

## **Product datasheet for MG227496**

### Ccr6 (NM 001190336) Mouse Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Ccr6 (NM\_001190336) Mouse Tagged ORF Clone

Tag: TurboGFP

Symbol: Ccr6

Synonyms: CC-CKR-6; CCR-6; Cmkbr6; KY411

Mammalian Cell

Selection:

Neomycin

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >MG227496 representing NM\_001190336 Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGAATTCCACAGAGTCCTACTTTGGAACGGATGATTATGACAACACAGAGTATTATTCTATTCCTCCAG ACCATGGGCCATGCTCCCTAGAAGAGGTCAGAAACTTCACCAAGGTATTTGTGCCAATTGCCTACTCCTT AATATGTGTCTTTGGCCTCCTGGGCAACATTATGGTGGTGATGACCTTTGCCTTCTACAAGAAAGCCAGA TCCATGACTGACGTCTACCTGTTGAACATGGCCATCACAGACATACTCTTTGTCCTCACCCTACCGTTCT GGGCAGTTACTCATGCCACCAACACTTGGGTTTTCAGCGATGCACTGTGTAAACTGATGAAAGGCACATA TGCGGTCAACTTTAACTGTGGGATGCTGCTCCTGGCCTGTATCAGCATGGACCGGTACATTGCCATCGTC CAGGCAACCAAATCTTTCCGGGTACGCTCCAGAACACTGACGCACAGTAAGGTCATCTGTGTGGCAGTGT GGTTCATCTCCATCATCTCAAGCCCTACATTTATCTTCAACAAGAAATACGAGCTGCAGGATCGTGA TGTCTGTGAGCCACGGTACAGGTCTGTCTCAGAGCCCATCACGTGGAAGCTGCTGGGTATGGGACTGGAG CTGTTCTTTGGGTTCTTCACCCCTTTGCTGTTTATGGTGTTCTGCTATCTGTTCATTATCAAGACCTTGG TGCAGGCCCAGAACTCCAAGAGGCACAGAGCCATCCGAGTCGTGATCGCTGTGGTTCTCGTGTTCCTGGC TTGTCAGATCCCTCACAACATGGTCCTCCTCGTGACTGCGGTCAACACGGGCAAAGTGGGCCGGAGCTGC AGCACCGAGAAAGTCCTCGCCTACACCAGGAACGTGGCCGAGGTCCTGGCTTTCCTGCATTGCTGCCTCA ACCCCGTGTTGTATGCGTTTATTGGACAGAAATTCAGAAACTACTTCATGAAGATCATGAAGGATGTGTG GTGTATGAGAAGGAAGAATAAGATGCCTGGCTTCCTCTGTGCCCGGGTTTACTCGGAAAGCTACATCTCC AGGCAGACCAGTGAGACCGTCGAAAATGATAATGCATCGTCCTTTACCATG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >MG227496 representing NM\_001190336

Red=Cloning site Green=Tags(s)


MNSTESYFGTDDYDNTEYYSIPPDHGPCSLEEVRNFTKVFVPIAYSLICVFGLLGNIMVVMTFAFYKKAR SMTDVYLLNMAITDILFVLTLPFWAVTHATNTWVFSDALCKLMKGTYAVNFNCGMLLLACISMDRYIAIV QATKSFRVRSRTLTHSKVICVAVWFISIIISSPTFIFNKKYELQDRDVCEPRYRSVSEPITWKLLGMGLE LFFGFFTPLLFMVFCYLFIIKTLVQAQNSKRHRAIRVVIAVVLVFLACQIPHNMVLLVTAVNTGKVGRSC STEKVLAYTRNVAEVLAFLHCCLNPVLYAFIGQKFRNYFMKIMKDVWCMRRKNKMPGFLCARVYSESYIS RQTSETVENDNASSFTM


TRTRPLE - GFP Tag - V

**Restriction Sites:** 

Sgfl-Mlul

**Cloning Scheme:** 





**ACCN:** NM\_001190336

ORF Size: 1101 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Cytogenetics:

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeq:** <u>NM 001190336.1</u>, <u>NP 001177265.1</u>

17 A1

 RefSeq Size:
 1324 bp

 RefSeq ORF:
 1104 bp

 Locus ID:
 12458


 UniProt ID:
 054689



#### **Gene Summary:**

Receptor for the C-C type chemokine CCL20. Binds to CCL20 and subsequently transduces a signal by increasing the intracellular calcium ion levels (PubMed:20068036). Although CCL20 is its major ligand it can also act as a receptor for non-chemokine ligands such as betadefensins (PubMed:25122636). Binds to defensin DEFB1 leading to increase in intracellular calcium ions and cAMP levels. Its binding to DEFB1 is essential for the function of DEFB1 in regulating sperm motility and bactericidal activity (By similarity). Binds to defensins DEFB4 and DEFB4A/B and mediates their chemotactic effects (PubMed:20068036). The ligandreceptor pair CCL20-CCR6 is responsible for the chemotaxis of dendritic cells (DC), effector/memory T-cells and B-cells and plays an important role at skin and mucosal surfaces under homeostatic and inflammatory conditions, as well as in pathology, including cancer and various autoimmune diseases. CCR6-mediated signals are essential for immune responses to microbes in the intestinal mucosa and in the modulation of inflammatory responses initiated by tissue insult and trauma (PubMed:21376174). CCR6 is essential for the recruitment of both the proinflammatory IL17 producing helper T-cells (Th17) and the regulatory T-cells (Treg) to sites of inflammation (PubMed:19050256). Required for the normal migration of Th17 cells in Peyers patches and other related tissue sites of the intestine and plays a role in regulating effector T-cell balance and distribution in inflamed intestine (PubMed:19129757). Plays an important role in the coordination of early thymocyte precursor migration events important for normal subsequent thymocyte precursor development, but is not required for the formation of normal thymic natural regulatory T-cells (nTregs). Required for optimal differentiation of DN2 and DN3 thymocyte precursors (PubMed:24638065). Essential for B-cell localization in the subepithelial dome of Peyers-patches and for efficient Bcell isotype switching to IgA in the Peyers-patches (PubMed:27174992). Essential for appropriate anatomical distribution of memory B-cells in the spleen and for the secondary recall response of memory B-cells (PubMed:25505290). Positively regulates sperm motility and chemotaxis via its binding to CCL20 (PubMed:23765988).[UniProtKB/Swiss-Prot Function]

# **Product images:**



Circular map for MG227496