

Product datasheet for **MG227569**

Mta1 (NM_054081) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Mta1 (NM_054081) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Mta1
Synonyms:	MGC118456
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MG227569 representing NM_054081
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCCGCCAACATGTACAGGGTCGGAGACTATGTCTACTTTGAGAACTCCTCCAGCAACCCGTACCTGA
 TCCGCAGGATTGAAGAGCTTAACAAGACAGCCAACGGGAATGTGGAGGCCAAAGTGGTGTGTTTCTACAG
 AAGGCGGGACATTTCCAGCAGCCTCATCGCGCTGGCGGACAAGCATGCAAGGGAAGTGGAGGAGGAGGTA
 GAAAACCCAGAGATGGTGGACCTGCCTGAGAACTTAAGCATCAGCTGCGGCATCGAGAAGTGTCTCTGT
 CTCGGCAGTTGGAGTCTCTGCCTGCCACCCACATCAGGGGCAAGTGCAGTGTACCCTGTCAATGAGAC
 GGAGTCACTCAAGTCTACTTGGAGCGTGGAGTTTCTTCTTCTACTCTAGTCTACGACCCACAGCAG
 AAGACCTCCTGGCTGATAAAGGGGAAATTCGTGTAGGAAACCGGTACCAGGCTGACATCACTGACTTGC
 TGAAAGAAGGTGAGGAGGACGGCCGCGATCAGTCAAACCTGGAGACCAAGGTGTGGGAAGCCACAATCC
 ACTTGTGGATAAGCAGATTGACCAGTTTCTGGTGGTGGCCGCTCTGTGGGTACCTTTGCACGGGCCCTG
 GATTGCAGCAGCTCCGTGCGGCAGCCTAGCCTGCACATGAGTGGCGCAGCGGCCTCCCGAGACATCACCC
 TGTTCATGCTATGGACACGCTGCACAAGAATCTACGACATCTCCAAGGCCATCTCGGCCCTCGTGCC
 CCAGGGTGGGCCAGTGTCTGCCGCGATGAGATGGAGGAGTGGTCCGCATCAGAAGCCAACCTTTTTGAG
 GAAGCTCTGGAATAATGGGAAAGATTTACCAGCATTCAACAAGATTTTCTCCATGGAAGTCGCTCA
 CCAGCATCATTGAGTATTACTACATGTGGAAGACCACCCGACAGATACGTCCAGCAGAAACGCTTGAAGC
 AGCTGAAGCTGAGAGCAAGTTAAGCAGGTTTATATCCCAACTATAACAAGCCCAACCCAAACCCAGATC
 AGCGCCAGCAGTGTCAAGGCCACTGTAGTGAATGGCACAGGGACACCAGGCCAGAGCCCCGGGGCCGGTC
 GGGCCTGCGAGAGCTGTTACACCACACAGTCTTACCAGTGGTATTCTTGGGGTCCCCTAACATGCAGTG
 TCGCCTCTGCGCATCTTGTGGACATATTGGAAGAAATATGGTGGCTTGAATAAGCCAAACCCGGTTAGAT
 GGAGAGAGGCCGGGACCAACCGCAATAACATGAGTCCCCATGGCATCCCAGCTCGGAGCAGTGGGAGCC
 CCAAGTTTGCCATGAAGACAAGGCAGGCCCTTCTACCTGCATACTACCAAGTTGACACGCATTGCCCGCG
 CTTGTGCCGTGAGATCCTACGCCATGGCATGCTGCCCGCACCCCTACATGCCATCAACAGCGCAGCA
 ATCAAGGCTGAATGCACAGCACGGCTGCCCGAAGCCTCCAGAGCCCACTGGTGTGAAGCAGGTAGTAC
 GGAAGCCCCTGGAGGCCGTGCTCCGGTATCTTGGAGCCATCCCCGCCCGCCTAAGCCCCGACCCCGTGA
 GAGTTCATCCAGTGTGCTCAGCAGTCTGACCCCTGCCAAGTCAGCCCTGTATCAACAACGGCTCCCCT
 ACCATCCTGGGCAAGAGGAGCTACGAGCAGCACAATGGGGTAGATGGCAACATGAAGAAGCGCCTTTGA
 TGCCCAAGTAGGGTCTGGCAAACCATGGACAGACCAGGCACATGGGACCAAGTCGGAATCTCCTGCTCAA
 TGGGAAGTCTACCCACCAAGTGCGCCTAATCCGTGGTGGCTCCCTGCCTCCAGTCAAGCGGGCGCGA
 ATGAACTGGATCGACGCCCCAGATGATGATTTTACATGGCCACCGAAGAGACCAGGAAAATCCGAAAGC
 TGCTCTCATCCTCAGAAACCAAGCGTGTGCCCGGCCCTACAACCCATTGCCCTGCGCCAGAGCCA
 GGCCCTGCCGCTGCGGCCACCCCACTGCACCAGTCAATGATGAGCCCATGTTATTGAGGAC

ACGCGTACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence: >MG227569 representing NM_054081
 Red=Cloning site Green=Tags(s)

MAANMYRVGDYVYFENSSSNPYLIRRIEELNKTANGNVEAKVVCFYRRRDISSSLIALADKHAREVEEEV
 ENPEMVDLPEKLRHRELFLSRQLESLPATHIRGKCSVTLLNETESLKSYLREDFFFYSLVYDPQQ
 KTLLADKGEIRVGNRYQADITDLLKEGEEDGRDQSKLETKVWEAHNPLVDKQIDQFLVVARSVGTFARAL
 DCSSSVRQPSLHMSAAAAARDITLFHAMDTLHKNIYDISKAISALVPQGGPVLCRDEMEEWSASEANLFE
 EALEKYGKDFTDIQQDFLPWKSLTSEIEYYMMWKTDRYVYQQRKLAEEAEKSKLKQVYIPNYNKPNNQI
 SASSVKATVVNGTGTGPGQSPGAGRACESCYTTQSYQWYWGPPNMQCRLCASCWTYWKYGLKMPTRL
 GERPGPNRNMSPHGIPARSSGSPKFMKTRQAFYLHTTKL TRIARRLCREILRPWHAARHPYMPINSAA
 IKAECTARLPEASQSPLVLKQVVRKPLEAVLRYLETHPRPPKPDVVKSSSVLSSLTPAKSAPVINNGSP
 TILGKRSYEQHNGVDGNMCKRLLMPSRGLANHGQTRHMGPSRNLNLLNGKSYPTKVRLIRGSSLPPVKRRR
 MNWIDAPDDVFYMATEETRKIRKLLSSSETKRAARRPKPIALRQSQUALPLRPPPPAPVNDPEIVIED



TRTRPLE - GFP Tag - V


Restriction Sites:

Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:


ACCN: NM_054081

ORF Size: 2094 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in *E. coli* are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_054081.2](#), [NP_473422.2](#)

RefSeq Size: 2775 bp

RefSeq ORF: 2097 bp

Locus ID: 116870

Cytogenetics: 12 F1

Gene Summary: Transcriptional coregulator which can act as both a transcriptional corepressor and coactivator. As a part of the histone-deacetylase multiprotein complex (NuRD), regulates transcription of its targets by modifying the acetylation status of the target chromatin and cofactor accessibility to the target DNA. In conjunction with other components of NuRD, acts as a transcriptional corepressor of BRCA1, ESR1, TFF1 and CDKN1A. Acts as a transcriptional coactivator of BCAS3, PAX5 and SUMO2, independent of the NuRD complex. Stimulates the expression of WNT1 by inhibiting the expression of its transcriptional corepressor SIX3. Regulates p53-dependent and -independent DNA repair processes following genotoxic stress. Regulates the stability and function of p53/TP53 by inhibiting its ubiquitination by COP1 and MDM2 thereby regulating the p53-dependent DNA repair. Plays an important role in tumorigenesis, tumor invasion, and metastasis. Plays a role in the regulation of the circadian clock and is essential for the generation and maintenance of circadian rhythms under constant light and for normal entrainment of behavior to light-dark (LD) cycles. Positively regulates the CLOCK-ARNTL/BMAL1 heterodimer mediated transcriptional activation of its own transcription and the transcription of CRY1. Regulates deacetylation of ARNTL/BMAL1 by regulating SIRT1 expression, resulting in derepressing CRY1-mediated transcription repression. With Tfcp2l1, promotes establishment and maintenance of pluripotency in embryonic stem cells (ESCs) and inhibits endoderm differentiation (PubMed:28982712). [UniProtKB/Swiss-Prot Function]