

Product datasheet for MR200006L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ost4 (NM_001134692) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Ost4 (NM 001134692) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ost4

Synonyms: 2310016E02Rik

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001134692

ORF Size: 114 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR200006).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001134692.1</u>, <u>NP 001128164.1</u>

 RefSeq Size:
 657 bp

 RefSeq ORF:
 114 bp

 Locus ID:
 67695

 UniProt ID:
 Q99LX8

 Cytogenetics:
 5 B1







Gene Summary:

Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity. Specifically involved in maintaining stability of STT3A-containing OST complexes.[UniProtKB/Swiss-Prot Function]