

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR200018L4V

Atp5e (NM_025983) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Atp5e (NM_025983) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Atp5e
Synonyms:	2410043G19Rik; ATPE; AV000645
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_025983
ORF Size:	159 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR200018).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 025983.3, NP 080259.1</u>
RefSeq Size:	419 bp
RefSeq ORF:	159 bp
Locus ID:	67126
UniProt ID:	<u>P56382</u>
Cytogenetics:	2 H4



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP
from ADP in the presence of a proton gradient across the membrane which is generated by
electron transport complexes of the respiratory chain. F-type ATPases consist of two
structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) -
containing the membrane proton channel, linked together by a central stalk and a peripheral
stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary
mechanism of the central stalk subunits to proton translocation. Part of the complex F(1)
domain and of the central stalk which is part of the complex rotary element. Rotation of the
central stalk against the surrounding alpha(3)beta(3) subunits leads to hydrolysis of ATP in
three separate catalytic sites on the beta subunits (By similarity).[UniProtKB/Swiss-Prot
Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US