

Product datasheet for **MR200081**

Gtf2h5 (NM_181392) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Gtf2h5 (NM_181392) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Gtf2h5
Synonyms: 2700017P07Rik; 2810432H05Rik; D17Wsu155e
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >MR200081 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGTCAACGTGCTGAAAGGGGTGCTTATAGAATGTGACCCTGCCATGAAGCAGTTCTTGCTGTACTTGG
ATGAGGCCAACGCCTTGGGGAAGAAGTTCATCATTCAGGACATTGATGACACGCACGTCTTCGT**CATTGC**
CGAGCTGGTCAACGTCTCCAGGAGCGAGTAGGGGAACTGATGGACCAGAATGCCTTTTCTT**ACCCAG**
AAG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR200081 protein sequence
Red=Cloning site Green=Tags(s)
MVNVLKGVLI~~ECDPAMKQFLLYLDEANALGKFKIIQDIDDTHVFVIAELVNVLQERVGELMDQNAFSLTQ~~
K

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI



[View online »](#)

Cloning Scheme:


ACCN: NM_181392

ORF Size: 216 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_181392.3](#)


RefSeq Size: 1191 bp

RefSeq ORF: 216 bp

Locus ID: 66467
UniProt ID: [Q8K2X8](#)
Cytogenetics: 17 3.7 cM
MW: 8 kDa

Gene Summary: Component of the general transcription and DNA repair factor IIH (TFIIH) core complex, which is involved in general and transcription-coupled nucleotide excision repair (NER) of damaged DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. In NER, TFIIH acts by opening DNA around the lesion to allow the excision of the damaged oligonucleotide and its replacement by a new DNA fragment. In transcription, TFIIH has an essential role in transcription initiation. When the pre-initiation complex (PIC) has been established, TFIIH is required for promoter opening and promoter escape. Phosphorylation of the C-terminal tail (CTD) of the largest subunit of RNA polymerase II by the kinase module CAK controls the initiation of transcription. Necessary for the stability of the TFIIH complex and for the presence of normal levels of TFIIH in the cell.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR200081