

## Product datasheet for **MR200291L4V**

### Cxcl10 (NM\_021274) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Cxcl10 (NM_021274) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Cxcl10
Synonyms:	C7; CRG-2; gIP-10; Ifi10; INP10; IP-10; IP10; mob-1; Scyb10
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_021274
ORF Size:	297 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR200291).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_021274.1</a>
RefSeq Size:	1120 bp
RefSeq ORF:	297 bp
Locus ID:	15945
UniProt ID:	<a href="#">P17515</a>
Cytogenetics:	5 46.57 cM



[View online »](#)

**Gene Summary:**

Pro-inflammatory cytokine that is involved in a wide variety of processes such as chemotaxis, differentiation, and activation of peripheral immune cells, regulation of cell growth, apoptosis and modulation of angiostatic effects (By similarity) (PubMed:28623423). Plays thereby an important role during viral infections by stimulating the activation and migration of immune cells to the infected sites (PubMed:18624292, PubMed:19017990, PubMed:28468883). Mechanistically, binding of CXCL10 to the CXCR3 receptor activates G protein-mediated signaling and results in downstream activation of phospholipase C-dependent pathway, an increase in intracellular calcium production and actin reorganization. In turn, recruitment of activated Th1 lymphocytes occurs at sites of inflammation (By similarity). Activation of the CXCL10/CXCR3 axis plays also an important role in neurons in response to brain injury for activating microglia, the resident macrophage population of the central nervous system, and directing them to the lesion site. This recruitment is an essential element for neuronal reorganization (PubMed:15456824).[UniProtKB/Swiss-Prot Function]