

## Product datasheet for **MR200317L3V**

### Sumo1 (NM\_009460) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Sumo1 (NM_009460) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Sumo1
Synonyms:	GMP1; PIC1; SENTRIN; SMT3; Smt3C; SMT3H3; SMTP3; SUMO-1; Ubl1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_009460
ORF Size:	306 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR200317).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_009460.2</a> , <a href="#">NP_033486.1</a>
RefSeq Size:	1230 bp
RefSeq ORF:	306 bp
Locus ID:	22218
UniProt ID:	<a href="#">P63166</a>
Cytogenetics:	1 C2



[View online »](#)

**Gene Summary:**

Ubiquitin-like protein that can be covalently attached to proteins as a monomer or a lysine-linked polymer. Covalent attachment via an isopeptide bond to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by E3 ligases such as PIAS1-4, RANBP2 or CBX4. This post-translational modification on lysine residues of proteins plays a crucial role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Involved for instance in targeting RANGAP1 to the nuclear pore complex protein RANBP2. Covalently attached to the voltage-gated potassium channel KCNB1; this modulates the gating characteristics of KCNB1. Polymeric SUMO1 chains are also susceptible to polyubiquitination which functions as a signal for proteasomal degradation of modified proteins. May also regulate a network of genes involved in palate development. Covalently attached to ZFH3 (By similarity).[UniProtKB/Swiss-Prot Function]