

## Product datasheet for MR200390L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Rbx1 (NM\_019712) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: Rbx1 (NM\_019712) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Rbx1

**Synonyms:** 1500002P15Rik; AA517855; ROC1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 019712

ORF Size: 327 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR200390).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 019712.3</u>, <u>NP 062686.1</u>

RefSeq Size: 1655 bp
RefSeq ORF: 327 bp
Locus ID: 56438
UniProt ID: P62878
Cytogenetics: 15 E1





## **Gene Summary:**

E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein ligase (CRLs) complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins, including proteins involved in cell cycle progression, signal transduction, transcription and transcription-coupled nucleotide excision repair (PubMed:22118460). CRLs complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins, ARIH1 mediating addition of the first ubiquitin on CRLs targets (By similarity). The functional specificity of the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition components (By similarity). As a component of the CSA complex promotes the ubiquitination of ERCC6 resulting in proteasomal degradation (By similarity). Through the RING-type zinc finger, seems to recruit the E2 ubiquitination enzyme, like CDC34, to the complex and brings it into close proximity to the substrate (By similarity). Probably also stimulates CDC34 autoubiquitination (By similarity). May be required for histone H3 and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair (By similarity). Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M (By similarity). Involved in the ubiquitination of KEAP1, ENC1 and KLHL41 (By similarity). In concert with ATF2 and CUL3, promotes degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM (By similarity).[UniProtKB/Swiss-Prot Function]