

# **Product datasheet for MR200405**

### Fkbp1a (NM 008019) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Fkbp1a (NM\_008019) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Fkbp1a

**Synonyms:** Fkb; Fkbp; Fkbp1; FKBP12

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR200405 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGGAGTGCAGGTGGAGACCATCTCTCCTGGAGACCGGGCCACCTTCCCAAAGCGCGGCCAGACCTGCG TGGTGCACTACACGGGGATGCTTGAAGATGGAAAGAAATTTGATTCCTCTCGGGACAGAAACAAGCCTTT TAAGTTTACACTAGGCAAGCAGGAGGTGATCCGAGGCTGGGAGGAAGGGGTAGCCCAGATGAGTGTGGGT CAGAGAGCCAAACTGATAATCTCCTCAGACTATGCCTATGGAGCCACCGGGCACCCAGGCATCATCCCAC

CACATGCCACTCTTGTTTTTGATGTGGAGCTTCTAAAACTGGAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >MR200405 protein sequence

Red=Cloning site Green=Tags(s)

MGVQVETISPGDGRTFPKRGQTCVVHYTGMLEDGKKFDSSRDRNKPFKFTLGKQEVIRGWEEGVAQMSVG

QRAKLIISSDYAYGATGHPGIIPPHATLVFDVELLKLE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_008019

ORF Size: 327 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



#### **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 008019.3</u>

 RefSeq Size:
 1667 bp

 RefSeq ORF:
 327 bp

 Locus ID:
 14225

 UniProt ID:
 P26883


 Cytogenetics:
 2 G3

**MW:** 11.9 kDa

**Gene Summary:** This gene is a member of the immunophilin family. The encoded protein is a cis-trans prolyl

isomerase that binds the immunosuppressants FK506 and rapamycin, and is associated with immunoregulation, protein folding, receptor signaling, protein trafficking and T-cell activation. It may modulate the calcium release activity of the ryanodine receptor Ryr1. It also interacts with the type I TGF-beta receptor. Disruption of this gene in mouse causes severe ventricular defects. Pseudogenes of this gene have been defined on chromosomes 4, 10, 14, and 16. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2014]

## **Product images:**



Circular map for MR200405