

## **Product datasheet for MR200451**

### Dad1 (NM\_010015) Mouse Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Dad1 (NM\_010015) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Dad1

Synonyms: Al323713

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR200451 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TTGCTGACTTCCTCTTTGCCAGCACGATCCTGCACCTTGTCGTCATGAACTTCGTTGGC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >MR200451 protein sequence

Red=Cloning site Green=Tags(s)

MSASVVSVISRFLEEYLSSTPQRLKLLDAYLLYILLTGALQFGYCLLVGTFPFNSFLSGFISCVGSFILA

VCLRIQINPQNKADFQGISPERAFADFLFASTILHLVVMNFVG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_010015

ORF Size: 342 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 010015.1</u>

RefSeq Size: 3506 bp RefSeq ORF: 342 bp



**Locus ID:** 13135 **UniProt ID:** P6180

UniProt ID: P61804

Cytogenetics: 14 27.7 cM

**MW:** 12.5 kDa

**Gene Summary:** Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a


defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-

pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent

polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs

cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity.[UniProtKB/Swiss-Prot Function]

# **Product images:**



Circular map for MR200451