

## **Product datasheet for MR200530**

## Eif4ebp1 (NM\_007918) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Eif4ebp1 (NM\_007918) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Eif4ebp1

Synonyms: 4e-bp1; AA959816; PHAS-I

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR200530 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

Т

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAG**GTTTAA** 

**Protein Sequence:** >MR200530 protein sequence

Red=Cloning site Green=Tags(s)

 ${\tt MSAGSSCSQTPSRAIPTRRVALGDGVQLPPGDYSTTPGGTLFSTTPGGTRIIYDRKFLMECRNSPVAKTP}$ 

PKDLPAIPGVTSPTSDEPPMQASQSQLPSSPEDKRAGGEESQFEMDI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_007918

ORF Size: 354 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 007918.3</u>, <u>NP 031944.3</u>

RefSeq Size: 981 bp RefSeq ORF: 354 bp Locus ID: 13685



 UniProt ID:
 Q60876

 Cytogenetics:
 8 15.95 cM


**MW:** 12.3 kDa

**Gene Summary:** Repressor of translation initiation that regulates EIF4E activity by preventing its assembly into

the eIF4F complex: hypophosphorylated form competes with EIF4G1/EIF4G3 and strongly binds to EIF4E, leading to repress translation. In contrast, hyperphosphorylated form dissociates from EIF4E, allowing interaction between EIF4G1/EIF4G3 and EIF4E, leading to initiation of translation (By similarity). Mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1

pathways (PubMed:7629182).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Circular map for MR200530