

## Product datasheet for MR200686L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Polr2f (NM\_027231) Mouse Tagged ORF Clone Lentiviral Particle

#### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Polr2f (NM\_027231) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Polr2f

**Synonyms:** 1810060D16Rik; RPB6

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 027231

ORF Size: 384 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR200686).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 027231.1, NP 081507.1</u>

 RefSeq Size:
 612 bp

 RefSeq ORF:
 384 bp

 Locus ID:
 69833

 UniProt ID:
 P61219

 Cytogenetics:
 15 37.7 cM







#### **Gene Summary:**

DNA-dependent RNA polymerases catalyze the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Common component of RNA polymerases I, II and III which synthesize ribosomal RNA precursors, mRNA precursors and many functional non-coding RNAs, and small RNAs, such as 5S rRNA and tRNAs, respectively. Pol II is the central component of the basal RNA polymerase II transcription machinery. Pols are composed of mobile elements that move relative to each other. In Pol II, POLR2F/RPB6 is part of the clamp element and together with parts of RPB1 and RPB2 forms a pocket to which the RPB4-RPB7 subcomplex binds (By similarity).[UniProtKB/Swiss-Prot Function]