

## Product datasheet for MR200694

## H2az1 (NM\_016750) Mouse Tagged ORF Clone

**Product data:** 

**Product Type: Expression Plasmids** 

**Product Name:** H2az1 (NM\_016750) Mouse Tagged ORF Clone

Tag: Myc-DDK

Synonyms: H2A.; H2A.Z; H2a.z-1; H2A.Z1; H2af; H2afz

H2az1

**Mammalian Cell** Neomycin

Selection:

**ORF Nucleotide** 

Symbol:

Vector: pCMV6-Entry (PS100001) E. coli Selection: Kanamycin (25 ug/mL) >MR200694 ORF sequence

Red=Cloning site Blue=ORF Green=Tags(s) Sequence:

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCTGGCGGTAAGGCTGGAAAGGACTCCGGAAAGGCCAAGACAAAGGCGGTTTCCCGCTCGCAGCGAG CCGGCTTGCAGTTCCCTGTGGGCCGTATTCATCGACACCTGAAATCTAGGACAACCAGCCACGGACGTGT GGGCGCGACCGCCGCTGTGTACAGCGCAGCCATCCTGGAGTACCTCACCGCAGAGGTACTTGAGTTGGCA GGAAATGCGTCAAAAGACTTAAAGGTAAAGCGTATCACCCCTCGTCACTTGCAGCTTGCTATACGTGGAG ATGAAGAATTGGATTCTCTGATCAAAGCTACCATTGCTGGTGGTGGTGTCATCCCACACATCCACAAATC

GCTGATCGGGAAGAAGACAACAGAAGACTGTT

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

>MR200694 protein sequence **Protein Sequence:** 

Red=Cloning site Green=Tags(s)

MAGGKAGKDSGKAKTKAVSRSQRAGLQFPVGRIHRHLKSRTTSHGRVGATAAVYSAAILEYLTAEVLELA

GNASKDLKVKRITPRHLQLAIRGDEELDSLIKATIAGGGVIPHIHKSLIGKKGQQKTV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_016750

ORF Size: 387 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 016750.3</u>, <u>NP 058030.1</u>


RefSeq Size: 1069 bp
RefSeq ORF: 387 bp
Locus ID: 51788
UniProt ID: P0C0S6
Cytogenetics: 3 G3

**MW:** 13.6 kDa

**Gene Summary:** 

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic development and indicate that lack of functional histone H2A leads to embryonic lethality. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2015]

## **Product images:**



Circular map for MR200694