

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for MR200694L4V

## H2afz (H2az1) (NM\_016750) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Lentiviral Particles
H2afz (H2az1) (NM_016750) Mouse Tagged ORF Clone Lentiviral Particle
H2az1
H2A.; H2A.Z; H2a.z-1; H2A.Z1; H2af; H2afz
Puromycin
pLenti-C-mGFP-P2A-Puro (PS100093)
mGFP
NM_016750
387 bp
The ORF insert of this clone is exactly the same as(MR200694).
The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<u>NM 016750.3, NP 058030.1</u>
1069 bp
387 bp
51788
<u>P0C0S6</u>
3 G3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



#### H2afz (H2az1) (NM\_016750) Mouse Tagged ORF Clone Lentiviral Particle – MR200694L4V

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic development and indicate that lack of functional histone H2A leads to embryonic lethality. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2015]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US