

## Product datasheet for MR201080L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Nudt16l1 (BC025569) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

Product Name: Nudt16l1 (BC025569) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Nudt16l1

**Synonyms:** 1110001K21Rik; 5330437l08Rik; Sdos

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 BC025569

 ORF Size:
 453 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR201080).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** BC025569, AAH25569

RefSeq Size: 1354 bp
RefSeq ORF: 455 bp
Locus ID: 66911

Cytogenetics: 16 2.48 cM







### **Gene Summary:**

Key regulator of TP53BP1 required to stabilize TP53BP1 and regulate its recruitment to chromatin. In absence of DNA damage, interacts with the tandem Tudor-like domain of TP53BP1, masking the region that binds histone H4 dimethylated at 'Lys-20' (H4K20me2), thereby preventing TP53BP1 recruitment to chromatin and maintaining TP53BP1 localization to the nucleus. Following DNA damage, ATM-induced phosphorylation of TP53BP1 and subsequent recruitment of RIF1 leads to dissociate NUDT16L1/TIRR from TP53BP1, unmasking the tandem Tudor-like domain and allowing recruitment of TP53BP1 to DNA double strand breaks (DSBs). Binds U8 snoRNA.[UniProtKB/Swiss-Prot Function]