

Product datasheet for MR201093L4

Oard1 (NM_207219) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Oard1 (NM_207219) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Oard1
Synonyms:	AI314976; AW558560
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR201093).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_207219
ORF Size:	459 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_207219.2
RefSeq Size:	1219 bp
RefSeq ORF:	459 bp
Locus ID:	106821
UniProt ID:	Q8R5F3
Cytogenetics:	17 C
Gene Summary:	<p>ADP-ribose glycohydrolase that hydrolyzes ADP-ribose and acts on different substrates, such as proteins ADP-ribosylated on glutamate and O-acetyl-ADP-D-ribose. Specifically acts as a glutamate mono-ADP-ribosylhydrolase by mediating the removal of mono-ADP-ribose attached to glutamate residues on proteins. Does not act on poly-ADP-ribosylated proteins: the poly-ADP-ribose chain of poly-ADP-ribosylated glutamate residues must be hydrolyzed into mono-ADP-ribosylated glutamate by PARG to become a substrate for OARD1. Deacetylates O-acetyl-ADP ribose, a signaling molecule generated by the deacetylation of acetylated lysine residues in histones and other proteins. Catalyzes the deacylation of O-acetyl-ADP-ribose, O-propionyl-ADP-ribose and O-butyryl-ADP-ribose, yielding ADP-ribose plus acetate, propionate and butyrate, respectively.[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MR201093L4