

## Product datasheet for MR201190L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cbx7 (NM\_144811) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: Cbx7 (NM 144811) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Cbx7

**Synonyms:** 1600014J01Rik; Al851678; D15Ertd417e

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_144811

ORF Size: 477 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(MR201190).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 144811.3, NP 659060.1

RefSeq Size: 2893 bp
RefSeq ORF: 477 bp
Locus ID: 52609
UniProt ID: Q8VDS3

Cytogenetics: 15 37.85 cM



## **Gene Summary:**

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:16537902, PubMed:22226355). PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Promotes histone H3 trimethylation at 'Lys-9' (H3K9me3) (By similarity). Binds to histone H3 trimethylated at 'Lys-9' (H3K9me3) or at 'Lys-27' (H3K27me3) (PubMed:16537902, PubMed:22226355). Trimethylation at 'Lys-27' (H3K27me3) is important for chromatin recruitment (PubMed:22226355, PubMed:16537902). May possibly also bind trimethylated lysine residues in other proteins (in vitro) (PubMed:16537902). Binds non-coding, single-stranded RNA and double-stranded RNA (PubMed:20541999, PubMed:16537902). Plays a role in the timely repression of differentiation-specific genes in pluripotent embryonic stem cells to maintain the undifferentiated state (PubMed:22226355). Regulator of cellular lifespan by maintaining the repression of CDKN2A, but not by inducing telomerase activity (PubMed:14647293).[UniProtKB/Swiss-Prot Function]