

Product datasheet for MR201211

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cdkn1a (NM 001111099) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Cdkn1a (NM_001111099) Mouse Tagged ORF Clone

Tag: Myc-DDK Symbol: Cdkn1a

Synonyms: CAP; CAP20; CDK; CDKI; Cdkn; Cdkn1; CI; CIP1; mda; mda6; P2; P21; p21C; p21Cip1; p21W;

p21WAF; SD; SDI1; Waf; Waf1

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) E. coli Selection: Kanamycin (25 ug/mL) **ORF Nucleotide** >MR201211 ORF sequence

Red=Cloning site Blue=ORF Green=Tags(s) Sequence:

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGTCCAATCCTGGTGATGTCCGACCTGTTCCGCACAGGAGCAAAGTGTGCCGTTGTCTCTTCGGTCCCG TGGACAGTGAGCAGTTGCGCCGTGATTGCGATGCGCTCATGGCGGGCTGTCTCCAGGAGGCCCGAGAACG GTGGAACTTTGACTTCGTCACGGAGACGCCGCTGGAGGGCAACTTCGTCTGGGAGCGCGTTCGGAGCCTA GGGCTGCCCAAGGTCTACCTGAGCCCTGGGTCCCGCAGCCGTGACGACCTGGGAGGGGACAAGAGGCCCA GTACTTCCTCTGCCCTGCTGCAGGGGCCAGCTCCGGAGGACCACGTGGCCTTGTCGCTGTCTTGCACTCT GGTGTCTGAGCGGCCTGAAGATTCCCCGGGTGGGCCCGGAACATCTCAGGGCCGAAAACGGAGGCAGACC

AGCCTGACAGATTTCTATCACTCCAAGCGCAGATTGGTCTTCTGCAAGAGAAAACCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR201211 protein sequence

Red=Cloning site Green=Tags(s)

MSNPGDVRPVPHRSKVCRCLFGPVDSEQLRRDCDALMAGCLQEARERWNFDFVTETPLEGNFVWERVRSL GLPKVYLSPGSRSRDDLGGDKRPSTSSALLQGPAPEDHVALSLSCTLVSERPEDSPGGPGTSQGRKRRQT

SLTDFYHSKRRLVFCKRKP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV







Restriction Sites:

Sgfl-Mlul

Cloning Scheme:

Cloning sites used for ORF Shuttling:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001111099

ORF Size: 480 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001111099.2

 RefSeq Size:
 1936 bp

 RefSeq ORF:
 480 bp

 Locus ID:
 12575

 UniProt ID:
 P39689




 Cytogenetics:
 17 15.12 cM

 MW:
 17.8 kDa

Gene Summary:

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at the G1 pahse. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]

Product images:



Circular map for MR201211