

Product datasheet for MR201267

Ubd (NM_023137) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Ubd (NM 023137) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Ubd

Synonyms: FAT10

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >MR201267 representing NM_023137

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCTTCTGTCCGCACCTGTGTTGTCCGTTCAGACCAATGGCGGTTAATGACCTTTGAGACCACTGAGA
ATGACAAAGTGAAGAAAATGAACATATTAGGTCCCAAACCAAGGTCTCTGTACAGGACCAGATCCT
TCTGCTAGACTCCAAAATCCTCAAGCCCCATCGAAAATTGTCATCCTATGGGATTGACAAGGAAACCACT
ATCCACCTTACCCTGAAGGTGGTGAAGCCCAGTGATGAAGAGCTGCCCTTGTTTCTGGTGGAGTCCAAAA
ACGAGGGGCAAAGGCACCTCCTCCGAGTTCGAAGATCCAGCTCAGTGGCCCAGGTGAAAGAGATGATCGA
GAGTGTGACCTCTGTGATCCCTAAGAAGCAGGTTGTGAATTGCAACGGAAAGAAGCTGGAAGATGGAAAG
ATCATGGCTGACTACAACATCAAGAGTGGCAGTTTGCTCTTTCTGACAACACACCTGCACTGGGGGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR201267 representing NM_023137

Red=Cloning site Green=Tags(s)

MASVRTCVVRSDQWRLMTFETTENDKVKKINEHIRSQTKVSVQDQILLLDSKILKPHRKLSSYGIDKETT IHLTLKVVKPSDEELPLFLVESKNEGQRHLLRVRRSSSVAQVKEMIESVTSVIPKKQVVNCNGKKLEDGK

IMADYNIKSGSLLFLTTHCTGG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_023137

ORF Size: 486 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 023137.3, NP 075626.1</u>

RefSeq Size: 1006 bp
RefSeq ORF: 489 bp
Locus ID: 24108
UniProt ID: P63072




Cytogenetics: 17 B1

MW: 18.8 kDa

Gene Summary: Ubiquitin-like protein modifier which can be covalently attached to target protein and

subsequently leads to their degradation by the 26S proteasome, in a NUB1-dependent manner. Probably functions as a survival factor. Promotes the expression of the proteasome subunit beta type-9 (PSMB9/LMP2). Regulates TNF-alpha-induced and LPS-mediated activation of the central mediator of innate immunity NF-kappa-B by promoting TNF-alpha-mediated proteasomal degradation of ubiquitinated-I-kappa-B-alpha. Required for TNF-alpha-induced p65 nuclear translocation in renal tubular epithelial cells (RTECs). May be involved in dendritic cell (DC) maturation, the process by which immature dendritic cells differentiate into fully competent antigen-presenting cells that initiate T-cell responses. Mediates mitotic non-disjunction and chromosome instability, in long-term in vitro culture and cancers, by abbreviating mitotic phase and impairing the kinetochore localization of MAD2L1 during the prometaphase stage of the cell cycle. May be involved in the formation of aggresomes when proteasome is saturated or impaired. Mediates apoptosis in a caspase-dependent manner, especially in renal epithelium and tubular cells during renal diseases.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201267