

Product datasheet for MR201284

Mafg (NM_010756) Mouse Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Mafg (NM_010756) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Mafg
Synonyms:	AA545192; C630022N07Rik
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR201284 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGCC</mark>
	ATGACGACCCCCAATAAAGGAAACAAGGCCTTAAAGGTGAAGCGGGAGCCAGGCGAGAATGGCACCAGCT TGACCGACGAGGAGCTGGTAACCATGTCGGTGCGAGAGTTGAACCAGCACCTGCGAGGCCTCTCCAAGGA AGAGATCATCCAGCTGAAGCAGCGGAGGCGCACACTGAAGAACCGGGGCTACGCGGGCCAGCTGCCGCGTC AAGCGGGTGACACAGAAGGAGGAGGAGCTGGAGAAGCAGAAGGCGGAGCTCCAGCAGGAGGTGGAGAAGCTGG CCTCGGAGAATGCCAGCATGAAGCTGGAGCTCGATGCCCTGCGGCTCCAAGTACGAGGCCCTGCAGAACTT TGCCAGGACCGTGGCCCGCAGCCTGTGGCCCCAGCTCGGGGTCCCCTTGCTGCTGGCCTGGGGCCCCTG GTTCCTGGCAAGGTGGCTGCACCAGCGTCACAAATAGTAAAGTCCAAGACGGATGCTCGGTCA
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG GTTTAA
Protein Sequence:	>MR201284 protein sequence <mark>Red</mark> =Cloning site Green=Tags(s)
	MTTPNKGNKALKVKREPGENGTSLTDEELVTMSVRELNQHLRGLSKEEIIQLKQRRRTLKNRGYAASCRV KRVTQKEELEKQKAELQQEVEKLASENASMKLELDALRSKYEALQNFARTVARSPVAPARGPLAAGLGPL VPGKVAATSVITIVKSKTDARS
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Restriction Sites:	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Cloning Scheme:




* The last codon before the Stop codon of the ORF

ACCN:	NM_010756
ORF Size:	489 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 010756.3, NP 034886.1</u>
RefSeq Size:	1720 bp
RefSeq ORF:	489 bp
Locus ID:	17134
UniProt ID:	<u>054790</u>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Mafg (NM_010756) Mouse Tagged ORF Clone – MR201284
Cytogenetics:	11 84.35 cM
MW:	17.9 kDa
Gene Summary:	Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves (PubMed:16738329, PubMed:9679061). However, they seem to serve as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins, such as NFE2, NFE2L1 and NFE2L2, and recruiting them to specific DNA-binding sites (PubMed:16738329, PubMed:9679061). Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NFE2L2 transcription factor. Transcription factor, component of erythroid-specific transcription factor NFE2L2. Activates globin gene expression when associated with NFE2L2 (By similarity). May be involved in signal transduction of extracellular H(+) (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201284

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US