

## Product datasheet for MR201284L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Mafg (NM\_010756) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Mafg (NM\_010756) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Mafg

**Synonyms:** AA545192; C630022N07Rik

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 010756

ORF Size: 489 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR201284).

•

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 010756.3, NP 034886.1

 RefSeq Size:
 1720 bp

 RefSeq ORF:
 489 bp

 Locus ID:
 17134

 UniProt ID:
 054790

Cytogenetics: 11 84.35 cM







## **Gene Summary:**

Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves (PubMed:16738329, PubMed:9679061). However, they seem to serve as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins, such as NFE2, NFE2L1 and NFE2L2, and recruiting them to specific DNA-binding sites (PubMed:16738329, PubMed:9679061). Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NFE2L2 transcription factor. Transcription factor, component of erythroid-specific transcription factor NFE2L2. Activates globin gene expression when associated with NFE2L2 (By similarity). May be involved in signal transduction of extracellular H(+) (By similarity). [UniProtKB/Swiss-Prot Function]