

## Product datasheet for MR201380L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Arpc4 (NM\_001170486) Mouse Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Arpc4 (NM\_001170486) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Arpc4

**Synonyms:** 20kDa; 5330419I20Rik; Al327076; p20-Arc

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001170486

ORF Size: 504 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR201380).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001170486.1

 RefSeq Size:
 2152 bp

 RefSeq ORF:
 237 bp

 Locus ID:
 68089

 UniProt ID:
 P59999

 Cytogenetics:
 6 E3







## **Gene Summary:**

Actin-binding component of the Arp2/3 complex, a multiprotein complex that mediates actin polymerization upon stimulation by nucleation-promoting factor (NPF). The Arp2/3 complex mediates the formation of branched actin networks in the cytoplasm, providing the force for cell motility. In addition to its role in the cytoplasmic cytoskeleton, the Arp2/3 complex also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA. The Arp2/3 complex promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs).[UniProtKB/Swiss-Prot Function]