

Product datasheet for **MR201391L3V**

Naa50 (NM_028108) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Naa50 (NM_028108) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Naa50
Synonyms:	2600005K24Rik; 2810441M03Rik; AW112078; Mak3; Mak3p; Nat5; Nat13; San
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_028108
ORF Size:	510 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR201391).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_028108.3
RefSeq Size:	4526 bp
RefSeq ORF:	507 bp
Locus ID:	72117
UniProt ID:	Q6PGB6
Cytogenetics:	16 B4



[View online »](#)

Gene Summary:

N-alpha-acetyltransferase that acetylates the N-terminus of proteins that retain their initiating methionine. Has a broad substrate specificity: able to acetylate the initiator methionine of most peptides, except for those with a proline in second position. Also displays N-epsilon-acetyltransferase activity by mediating acetylation of the side chain of specific lysines on proteins. Autoacetylates in vivo. The relevance of N-epsilon-acetyltransferase activity is however unclear: able to acetylate H4 in vitro, but this result has not been confirmed in vivo. Component of a N-alpha-acetyltransferase complex containing NAA10 and NAA15, but NAA50 does not influence the acetyltransferase activity of NAA10: this multiprotein complex probably constitutes the major contributor for N-terminal acetylation at the ribosome exit tunnel, with NAA10 acetylating all amino termini that are devoid of methionine and NAA50 acetylating other peptides. Required for sister chromatid cohesion during mitosis by promoting binding of CDCA5/sororin to cohesin: may act by counteracting the function of NAA10.[UniProtKB/Swiss-Prot Function]