

Product datasheet for MR201427

Nudt15 (NM 172527) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Nudt15 (NM_172527) Mouse Tagged ORF Clone

Tag: Myc-DDK Symbol: Nudt15

Synonyms: 6530403O17; A730068G11Rik; MTH2

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR201427 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

AAAGGACCACGAAGACTCCT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR201427 protein sequence

Red=Cloning site Green=Tags(s)

MAANAEPRRRPGVGVGVVVLSCEHPRCVLLGKRKGSFGAGSFQLPGGHLEFGETWEECAQRETWEEAGLH LKNVCFASVVNSFVEKENYHYVTILMKGEVDMTHDSEPRNMEPEKNESWEWVPWEEFPPLDQLFWALRCL

KEQGYDPFKEDLNHLEGYRGEHLERTTKTP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM 172527

ORF Size: 513 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 172527.3

 RefSeq Size:
 2592 bp

 RefSeq ORF:
 513 bp

 Locus ID:
 214254

 UniProt ID:
 Q8BG93

 Cytogenetics:
 14 D3




MW:

19.6 kDa

Gene Summary:

May catalyze the hydrolysis of nucleoside triphosphates including dGTP, dTTP, dCTP, their oxidized forms like 8-oxo-dGTP and the prodrug thiopurine derivatives 6-thio-dGTP and 6-thio-GTP (PubMed:12767940). Could also catalyze the hydrolysis of some nucleoside diphosphate derivatives (By similarity). Hydrolyzes oxidized nucleosides triphosphates like 8-oxo-dGTP in vitro, but the specificity and efficiency towards these substrates are low. Therefore, the potential in vivo sanitizing role of this enzyme, that would consist in removing oxidatively damaged forms of nucleosides to prevent their incorporation into DNA, is unclear (PubMed:12767940). Through the hydrolysis of thioguanosine triphosphates may participate in the catabolism of thiopurine drugs (By similarity). May also have a role in DNA synthesis and cell cycle progression by stabilizing PCNA (By similarity). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201427