

Product datasheet for MR201427L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nudt15 (NM_172527) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nudt15 (NM 172527) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Nudt15

Synonyms: 6530403O17; A730068G11Rik; MTH2

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 172527

Tag: Myc-DDK

ORF Size: 513 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR201427).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 172527.1</u>

 RefSeq Size:
 2592 bp

 RefSeq ORF:
 513 bp

 Locus ID:
 214254

 UniProt ID:
 Q8BG93

Cytogenetics: 14 D3







Gene Summary:

May catalyze the hydrolysis of nucleoside triphosphates including dGTP, dTTP, dCTP, their oxidized forms like 8-oxo-dGTP and the prodrug thiopurine derivatives 6-thio-dGTP and 6-thio-GTP (PubMed:12767940). Could also catalyze the hydrolysis of some nucleoside diphosphate derivatives (By similarity). Hydrolyzes oxidized nucleosides triphosphates like 8-oxo-dGTP in vitro, but the specificity and efficiency towards these substrates are low. Therefore, the potential in vivo sanitizing role of this enzyme, that would consist in removing oxidatively damaged forms of nucleosides to prevent their incorporation into DNA, is unclear (PubMed:12767940). Through the hydrolysis of thioguanosine triphosphates may participate in the catabolism of thiopurine drugs (By similarity). May also have a role in DNA synthesis and cell cycle progression by stabilizing PCNA (By similarity). [UniProtKB/Swiss-Prot Function]