

# Product datasheet for MR201451L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Polr2g (NM\_026329) Mouse Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** Polr2g (NM\_026329) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Polr2g

**Synonyms:** 2410046K11Rik; A230108L04Rik; C76415; RBP7; Rpo2-7l

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 026329

Tag: Myc-DDK

ORF Size: 519 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR201451).

Sequence:

ACCN:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 026329.1, NP 080605.1

 RefSeq Size:
 868 bp

 RefSeq ORF:
 519 bp

 Locus ID:
 67710

 UniProt ID:
 P62488

Cytogenetics: 19 A







#### **Gene Summary:**

DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Component of RNA polymerase II which synthesizes mRNA precursors and many functional non-coding RNAs. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. RPB7 is part of a subcomplex with RPB4 that binds to a pocket formed by RPB1, RPB2 and RPB6 at the base of the clamp element. The RBP4-RPB7 subcomplex seems to lock the clamp via RPB7 in the closed conformation thus preventing double-stranded DNA to enter the active site cleft. The RPB4-RPB7 subcomplex binds single-stranded DNA and RNA. Binds RNA (By similarity).[UniProtKB/Swiss-Prot Function]