

Product datasheet for **MR201468L4V**

Pdpn (NM_010329) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Pdpn (NM_010329) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Pdpn
Synonyms:	Gp38; OTS-8; RANDAM-2; T1-alpha; T1a; T1alpha
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_010329
ORF Size:	519 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR201468).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_010329.2
RefSeq Size:	1839 bp
RefSeq ORF:	519 bp
Locus ID:	14726
UniProt ID:	Q62011
Cytogenetics:	4 E1



[View online »](#)

Gene Summary:

Mediates effects on cell migration and adhesion through its different partners. During development plays a role in blood and lymphatic vessels separation by binding CLEC1B, triggering CLEC1B activation in platelets and leading to platelet activation and/or aggregation (PubMed:14522983, PubMed:15231832, PubMed:20110424, PubMed:17616532). Interaction with CD9, on the contrary, attenuates platelet aggregation and pulmonary metastasis induced by PDPN. Mediates effects on cell migration and adhesion through its different partners. Through MSN or EZR interaction promotes epithelial-mesenchymal transition (EMT) leading to ERZ phosphorylation and triggering RHOA activation leading to cell migration increase and invasiveness. Interaction with CD44 promotes directional cell migration in epithelial and tumor cells (By similarity). In lymph nodes (LNs), controls fibroblastic reticular cells (FRCs) adhesion to the extracellular matrix (ECM) and contraction of the actomyosin by maintaining ERM proteins (EZR; MSN and RDX) and MYL9 activation through association with unknown transmembrane proteins. Engagement of CLEC1B by PDPN promotes FRCs relaxation by blocking lateral membrane interactions leading to reduction of ERM proteins (EZR; MSN and RDX) and MYL9 activation (PubMed:25347465). Through binding with LGALS8 may participate to connection of the lymphatic endothelium to the surrounding extracellular matrix (By similarity). In keratinocytes, induces changes in cell morphology showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion (PubMed:10574709). Controls invadopodia stability and maturation leading to efficient degradation of the extracellular matrix (ECM) in tumor cells through modulation of RHOC activity in order to activate ROCK1/ROCK2 and LIMK1/LIMK2 and inactivation of CFL1 (By similarity). Required for normal lung cell proliferation and alveolus formation at birth (PubMed:12654292). Does not function as a water channel or as a regulator of aquaporin-type water channels (By similarity). Does not have any effect on folic acid or amino acid transport (PubMed:12032185).[UniProtKB/Swiss-Prot Function]