

Product datasheet for MR201576

Rpl11 (NM_025919) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Rpl11 (NM_025919) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Rpl11

Synonyms: 2010203J19Rik

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >MR201576 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCGCAAGATCAAGGGGAAAAGGAGAACCCCATGCGGGAACTGCGCATCCGCAAGCTCTGCCTCAATA
TCTGCGTCGGGGAGAGCGGAGACAGACTGACCCGGGCAGCCAAGGTGTTGGAGCAGCTCACAGGCCAGAC
CCCGGTGTTCTCCAAAGCTAGATACACTGTCAGGTCCTTTGGCATCCGGAGAAATGAGAAGATTGCTGTT
CACTGCACAGTCCGCGAGCCAAGGCAGAGGAAATTCTGGAGAAAGGCCTGAAGGTGCGGGAGTATGAGT
TGCGGAAAAATAACTTCTCGGATACTGGAAACTTTGGTTTTTGGAATTCAAGAACACATTGACCTGGGCAT
CAAATACGACCCAAGCATTGGGATCTACGGCCTGGACTTCTATGTGGTGCTGGGTAGGCCAGGGTTCAGC
ATCGCAGACAAGAAGCGCAGAACAGGCTGCATTGGGGCCAAACACAGAATCAGCAAGGAGGAGGAGGCCATGC

GCTGGTTCCAGCAGAAGTACGATGGAATCATCCTTCCTGGAAAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR201576 protein sequence

Red=Cloning site Green=Tags(s)

MAQDQGEKENPMRELRIRKLCLNICVGESGDRLTRAAKVLEQLTGQTPVFSKARYTVRSFGIRRNEKIAV HCTVRGAKAEEILEKGLKVREYELRKNNFSDTGNFGFGIQEHIDLGIKYDPSIGIYGLDFYVVLGRPGFS

IADKKRRTGCIGAKHRISKEEAMRWFQQKYDGIILPGK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM 025919

ORF Size: 537 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 025919.3</u>

RefSeq Size: 594 bp
RefSeq ORF: 537 bp
Locus ID: 67025
UniProt ID: Q9CXW4
Cytogenetics: 4 D3




MW:

20.3 kDa

Gene Summary:

Component of the ribosome, a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell. The small ribosomal subunit (SSU) binds messenger RNAs (mRNAs) and translates the encoded message by selecting cognate aminoacyl-transfer RNA (tRNA) molecules. The large subunit (LSU) contains the ribosomal catalytic site termed the peptidyl transferase center (PTC), which catalyzes the formation of peptide bonds, thereby polymerizing the amino acids delivered by tRNAs into a polypeptide chain. The nascent polypeptides leave the ribosome through a tunnel in the LSU and interact with protein factors that function in enzymatic processing, targeting, and the membrane insertion of nascent chains at the exit of the ribosomal tunnel. As part of the 5S RNP/5S ribonucleoprotein particle it is an essential component of the LSU, required for its formation and the maturation of rRNAs. It also couples ribosome biogenesis to p53/TP53 activation. As part of the 5S RNP it accumulates in the nucleoplasm and inhibits MDM2, when ribosome biogenesis is perturbed, mediating the stabilization and the activation of TP53 (PubMed:21804542). Promotes nucleolar location of PML (PubMed:15195100).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201576