

Product datasheet for MR201636

Arf1 (NM 001130408) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Arf1 (NM_001130408) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Arf1

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >MR201636 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGGGAATATCTTTGCAAACCTCTTCAAGGGCCTTTTTGGCAAAAAGAAATGCGCATTCTCATGGTGG
GCCTGGATGCTGCAGGGAAGACAACAATTCTATACAAACTTAAGCTGGGCGAAATTGTGACCACCATTCC
CACCATTGGTTTCAATGTGGAGACTGTTGAATACAAGAATATCAGCTTCACCGTGTGGGATGTGGGCGGC
CAGGACAAGATCCGGCCGCTGTGGCGCCACTACTTCCAGAACACCCAAGGCTTGATCTTCGTAGTGGACA
GCAATGACAGAGAGCGTGTGAACGAGGCCCGTGAAGAGCTCATGAGGATGCTAGCTGAAGATGACCCG
AGATGCTGTTCTCTTGGTGTTTGCCAACAAGCAGGACCTCCCCAATGCCATGAATGCGGCCGAAATCACA
GACAAGCTGGGGCTGCACTCTCTACGCCACAGGAACTGGTACATTCAGGCCACCTGTGCCACCAGCGGGG

ACGGGCTCTATGAAGGACTAGATTGGCTGTCTAATCAGCTCCGGAACCAGAAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR201636 protein sequence

Red=Cloning site Green=Tags(s)

MGNIFANLFKGLFGKKEMRILMVGLDAAGKTTILYKLKLGEIVTTIPTIGFNVETVEYKNISFTVWDVGG QDKIRPLWRHYFQNTQGLIFVVDSNDRERVNEAREELMRMLAEDELRDAVLLVFANKQDLPNAMNAAEIT

DKLGLHSLRHRNWYIQATCATSGDGLYEGLDWLSNQLRNQK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM 001130408

ORF Size: 543 bp

OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.



Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 001130408.1</u>, <u>NP 001123880.1</u>

 RefSeq Size:
 1991 bp

 RefSeq ORF:
 546 bp

 Locus ID:
 11840

 UniProt ID:
 P84078


 Cytogenetics:
 11 B1.3

 MW:
 20.7 kDa

Gene Summary: GTP-binding protein involved in protein trafficking among different compartments

(PubMed:11950392). Modulates vesicle budding and uncoating within the Golgi complex. Deactivation induces the redistribution of the entire Golgi complex to the endoplasmic reticulum, suggesting a crucial role in protein trafficking. In its GTP-bound form, its triggers the association with coat proteins with the Golgi membrane. The hydrolysis of ARF1-bound GTP, which is mediated by ARFGAPs proteins, is required for dissociation of coat proteins from Golgi membranes and vesicles. The GTP-bound form interacts with PICK1 to limit PICK1-mediated inhibition of Arp2/3 complex activity; the function is linked to AMPA receptor (AMPAR) trafficking, regulation of synaptic plasicity of excitatory synapses and spine shrinkage during long-term depression (LTD).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201636