

Product datasheet for **MR201775L3**

Myo3b (BC034907) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: Myo3b (BC034907) Mouse Tagged Lenti ORF Clone
 Tag: Myc-DDK
 Symbol: Myo3b
 Synonyms: A430065P19Rik
 Mammalian Cell Selection: Puromycin
 Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)
 E. coli Selection: Chloramphenicol (34 ug/mL)
 ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as(MR201775).
 Restriction Sites: SgfI-MluI
 Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN: BC034907
 ORF Size: 564 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC034907.1
RefSeq Size:	1363 bp
RefSeq ORF:	566 bp
Locus ID:	329421
Cytogenetics:	2 C2
Gene Summary:	Probable actin-based motor with a protein kinase activity (By similarity). Required for normal cochlear hair bundle development and hearing. Plays an important role in the early steps of cochlear hair bundle morphogenesis. Influences the number and lengths of stereocilia to be produced and limits the growth of microvilli within the forming auditory hair bundles thereby contributing to the architecture of the hair bundle, including its staircase pattern (PubMed:26754646). Involved in the elongation of actin in stereocilia tips by transporting the actin regulatory factor ESPN to the plus ends of actin filaments (PubMed:22264607). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201775L3