

Product datasheet for MR201830

Cdc42 (NM_009861) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Cdc42 (NM_009861) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Cdc42
Synonyms: A1747189; AU018915
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >MR201830 representing NM_009861
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGCAGACAATTAAGTGTGTTGTTGGTGGTGGTGGTAAAACATGTCTCCTGATATCCTACA
 CAACAAACAAATCCCATCGGAATATGTACCAACTGTTTTGACAACTATGCAGTCACAGTTATGATTGG
 TGGAGAGCCATACACTCTGGACTTTTTGATACTGCAGGGCAAGAGGATTATGACGGACTACGACCCTA
 AGTTATCCACAGACAGATGTTTTCTAGTATGTTCTCAGTGGTCTCTCCATCCTATTTGAAAAATGTA
 AAGAAAAGTGGTGCCTGAGATAACTCACCCTGTCCAAAGACTCTTTCTTGCTTGTGGACCCAAAT
 TGATCTCAGAGATGACCCCTCTACTATTGAGAACTTGCCAAGAACAACAGAAGCCTATTACTCCAGAG
 ACTGCTGAAAAGCTGGCGCGGATCTGAAGCTGTCAAGTATGTGGAGTGTCTGCCCTCACACAGAAAG
 GCCTAAAGAATGTTTGTGGAGCAATATTGGCTGCCCTGGAGCCTCCAGAACCGAAGAAGAGCCGAG
 GTGTGTGCTGCTA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR201830 representing NM_009861
 Red=Cloning site Green=Tags(s)

MQTIKCVVVDGAVGKTCLLISYTTNKFPEYVPTVFDNYAVTVMIGGEPYTLGLFDTAGQEDYDGLRPL
 SYPQTDVFLVCFVSVSPSSFENVKEKWPEITHHCPKTPFLLVGTQIDLRDDPSTIEKLAKNKQKPIPE
 TAEKLARDLKAVKYVECSALTQKGLKNVFDGAILAALEPPPEPKSRRCVLL


TRTRPLEQKLISEEDLAANDILDYKDDDDKV



[View online »](#)

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_009861

ORF Size: 573 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_009861.1](#), [NM_009861.2](#), [NM_009861.3](#), [NP_033991.1](#)

RefSeq Size: 2063 bp

RefSeq ORF: 576 bp

Locus ID: 12540


UniProt ID: [P60766](#)

Cytogenetics: 4 69.83 cM

MW: 21.7 kDa

Gene Summary: Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state (PubMed:24352656). In its active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase. Regulates cell migration (By similarity). In neurons, plays a role in the extension and maintenance of the formation of filopodia, thin and actin-rich surface projections. Required for DOCK10-mediated spine formation in Purkinje cells and hippocampal neurons (PubMed:25851601). Facilitates filopodia formation upon DOCK11-activation (PubMed:22494997). Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Also plays a role in phagocytosis through organization of the F-actin cytoskeleton associated with forming phagocytic cups (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201830