

Product datasheet for MR201866L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cldn10 (NM_001160097) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cldn10 (NM_001160097) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Cldn10

Synonyms: 6720456l16Rik; Cldn; Cldn10a; Cldn10b; D14Ertd728; D14Ertd728e

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001160097

ORF Size: 582 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR201866).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001160097.1, NP 001153569.1

 RefSeq Size:
 1092 bp

 RefSeq ORF:
 582 bp

 Locus ID:
 58187

 UniProt ID:
 Q9Z0S6

Cytogenetics: 14 62.55 cM







Gene Summary:

This intronless gene encodes a member of the claudin family. Claudins are integral membrane proteins and components of tight unction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. Six alternatively spliced transcript variants have been identified, which encode different isoforms with distinct electric charge of the first extracellular loop and with or without the fourth transmembrane region. These isoforms exhibit distinct localization and function in paracellular anion or cation permeability. [provided by RefSeq, Aug 2010]