

## **Product datasheet for MR201957L3V**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cdkn1b (NM\_009875) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: Cdkn1b (NM 009875) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Cdkn1b

**Synonyms:** AA408329; AI843786; Kip1; p27; p27Kip1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 009875

ORF Size: 594 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR201957).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 009875.2</u>

 RefSeq Size:
 2405 bp

 RefSeq ORF:
 594 bp

 Locus ID:
 12576

 UniProt ID:
 P46414

 Cytogenetics:
 6 65.77 cM







## **Gene Summary:**

Important regulator of cell cycle progression (PubMed:8033213, PubMed:12972555). Inhibits the kinase activity of CDK2 bound to cyclin A, but has little inhibitory activity on CDK2 bound to SPDYA (By similarity). Involved in G1 arrest. Potent inhibitor of cyclin E- and cyclin A-CDK2 complexes (PubMed:8033213). Forms a complex with cyclin type D-CDK4 complexes and is involved in the assembly, stability, and modulation of CCND1-CDK4 complex activation. Acts either as an inhibitor or an activator of cyclin type D-CDK4 complexes depending on its phosphorylation state and/or stoichometry.[UniProtKB/Swiss-Prot Function]