

## Product datasheet for **MR202051L4V**

### **Psmb2 (NM\_011970) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Psmb2 (NM_011970) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Psmb2
Synonyms:	AU045357; AW108089; C7-I; D4Wsu33e; HC7-I
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_011970
ORF Size:	606 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR202051).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_011970.2</a> , <a href="#">NP_036100.3</a>
RefSeq Size:	861 bp
RefSeq ORF:	606 bp
Locus ID:	26445
UniProt ID:	<a href="#">Q9R1P3</a>
Cytogenetics:	4 60.9 cM



[View online »](#)

**Gene Summary:**

Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).[UniProtKB/Swiss-Prot Function]