

Product datasheet for MR202075

Rab13 (NM_026677) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Rab13 (NM_026677) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Rab13
Synonyms: 0610007N03Rik; B230212B15Rik
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >MR202075 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCCAAAGCCTACGACCACCTCTTCAAGTTGCTGCTCATCGGGGACTCGGGGGTGGCAAGACTTGTC
 TGATCATTCGCTTTCAGAGGACAACCTCAACAGCACTTACATCTCTACCATCGGAATTGATTTCAAGAT
 CCGAACCGTGGACATAGAGGGGAAGAGGATCAAAGTCAAGTGTGGGACACGGCTGGCCAAGAACGATTC
 AAGACAATAACTACCGCCTATTACCGTGGAGCCATGGGCATTATCCTCGTATATGACATCACAGATGAGA
 AATCCTTCGAGAATATTCAGAACTGGATGAAAAGCATCAAAGAGAATGCCTCTGCGGGAGTGGAGCGCCT
 CCTGCTGGGAAACAAGTGTGACATGGAGGCCAAGCGGCAGGTGCAGAGAGAGCAGGCGGAGAAGTTGGCT
 CGAGAGCACAGAATCCGATTTTTGAGACGAGTGCCAAATCCAGTGTGAATGTGGATGAGGCTTTAGTT
 CCCTGGCCCGTGACATCTTGCTCAAGACAGGAGCCGGAGATCGGGAACCAACAGTAAGCCCTCAAGCAC
 TGGCCTGAAAACATCTGACAAGAAGAACAAGTGTCTTGTAGGC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR202075 protein sequence
 Red=Cloning site Green=Tags(s)

MAKAYDHLFKLLIGDSGVGKTCIIIRFAEDNFNSTYISTIGIDFKIRTVDIEGKRIKLQVWDTAGQERF
 KTITTAYYRGAMGIILVYDITDEKSFENIQNWMKSIKENASAGVERLLLGNKCDMEAKRQVQREQAELKLA
 REHRIRFFETSASKSSVNVDEAFSSLARDILLKTGGRRSGTNSKPSSTGLKTSDDKKNKCLLG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



[View online »](#)

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_026677

ORF Size: 609 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_026677.4](#)

RefSeq Size: 1548 bp

RefSeq ORF: 609 bp

Locus ID: 68328


UniProt ID: [Q9DD03](#)

Cytogenetics: 3 39.21 cM

MW: 22.8 kDa

Gene Summary: The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in endocytic recycling and regulates the transport to the plasma membrane of transmembrane proteins like the tight junction protein OCLN/occludin. Thereby, it regulates the assembly and the activity of tight junctions. Moreover, it may also regulate tight junction assembly by activating the PKA signaling pathway and by reorganizing the actin cytoskeleton through the activation of the downstream effectors PRKACA and MICALL2 respectively. Through its role in tight junction assembly, may play a role in the establishment of Sertoli cell barrier. Plays also a role in angiogenesis through regulation of endothelial cells chemotaxis. Also involved in neurite outgrowth. Has also been proposed to play a role in post-Golgi membrane trafficking from the TGN to the recycling endosome. Finally, it has been involved in insulin-induced transport to the plasma membrane of the glucose transporter GLUT4 and therefore may play a role in glucose homeostasis.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR202075